



# Slower-X: reduced efficiency of selection in the early stages of X chromosome evolution

Andrea Mrnjavac, Ksenia A. Khudiakova, Nicholas H. Barton,  Beatriz Vicoso 

Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

Corresponding authors: Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria. Email: [andrea.mrnjavac@ist.ac.at](mailto:andrea.mrnjavac@ist.ac.at); Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria. Email: [bvicoso@ist.ac.at](mailto:bvicoso@ist.ac.at)

## Abstract

Differentiated X chromosomes are expected to have higher rates of adaptive divergence than autosomes, if new beneficial mutations are recessive (the “faster-X effect”), largely because these mutations are immediately exposed to selection in males. The evolution of X chromosomes after they stop recombining in males, but before they become hemizygous, has not been well explored theoretically. We use the diffusion approximation to infer substitution rates of beneficial and deleterious mutations under such a scenario. Our results show that selection is less efficient on diploid X loci than on autosomal and hemizygous X loci under a wide range of parameters. This “slower-X” effect is stronger for genes affecting primarily (or only) male fitness, and for sexually antagonistic genes. These unusual dynamics suggest that some of the peculiar features of X chromosomes, such as the differential accumulation of genes with sex-specific functions, may start arising earlier than previously appreciated.

**Keywords:** sheltering, faster-X theory, X chromosome, young sex chromosomes, degeneration, adaptation, demasculinization

## Layman Summary

In many species sex is determined by a pair of sex chromosomes, usually termed X and Y. Humans share sex chromosomes with all mammals apart from monotremes like platypus and echidna, but sex chromosomes evolved independently from a pair of autosomes many times in evolution. Sex chromosome evolution usually involves the degeneration of the Y chromosome, which loses most of the ancestral gene content. For example, the human Y chromosome contains around 80 genes, compared to 800 on the X. This means that XY males carry the majority of their X-linked genes in only one copy (as opposed to two copies of genes that are on the autosomes). So far, studies of sex chromosome evolution focused on evolutionarily old X chromosomes with degenerated Y chromosomes. This is in part because old sex chromosomes are easier to identify due to the extensive differences between X and Y. With the increasing availability of sequencing, large-scale population genomic studies have detected unusual patterns of evolution on young X chromosomes, but the evolutionary forces at play are currently unknown. Here, we model the evolutionary dynamics of young X chromosomes with a non-degenerated Y and show that young X chromosomes can experience degeneration, and demasculinization, that is, loss of male-important genes. This is in contrast with the prevailing perspective that X chromosomes with non-degenerated Y counterparts have similar evolutionary dynamics to autosomes.

## Introduction

In many species, the sex of an individual is determined by a pair of sex chromosomes, such as the X and Y of mammals, or Z and W in the case of the female heterogametic system (Bachtrog et al., 2014). Although we focus on the more commonly studied XY case, the models discussed here also apply to ZW systems (by switching the sexes). Sex chromosomes arise from a pair of autosomes when one of them acquires a sex-determining gene (Vicoso, 2019; Wright et al., 2016). This is often coupled with the suppression of recombination between X and Y chromosomes in the region surrounding the sex-determining gene. Recombination suppression can spread progressively along the chromosomes in a step-wise manner, creating distinct “evolutionary strata,” i.e., regions that stopped recombining at the same time, and which consequently have different levels of XY divergence (Jefferies et al., 2021; Ponnikas et al., 2018). Recombination suppression between

X and the Y reduces the efficiency of selection on the Y chromosome, leading to progressive gene loss on the Y (Bachtrog, 2013; Engelstädter, 2008). The resulting imbalance in gene copy number in males drives the evolution of dosage compensation mechanisms, which results in equal expression levels in males and females in somatic tissues. In male gonads, on the other hand, X chromosomes are often downregulated or completely inactivated (Larson et al., 2018; Mahadevaraju et al., 2021). In addition to this unusual regulatory architecture, X chromosomes have been found to differ from autosomes in various ways. One consistent feature is the over- and under-representation of genes with sex-specific patterns of expression (sex-biased genes), although the specific direction of the enrichment varies across species. For instance, the *Drosophila* X chromosome has a deficit of male-biased genes, whereas the mammalian X is enriched for genes with male-specific functions (Gurbich & Bachtrog, 2008). X chromosomes also

Received October 12, 2022; revisions received November 8, 2022; accepted December 17, 2022

© The Author(s) 2023. Published by Oxford University Press on behalf of The Society for the Study of Evolution (SSE) and European Society for Evolutionary Biology (ESEB).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

often have more transposable elements and repeats and have different gene densities, than autosomes. Finally, genes move out of X chromosomes more often than expected in both mammals and *Drosophila* (Gurbich & Bachtrog, 2008). Understanding what evolutionary processes drive these patterns has been the goal of extensive theoretical and empirical research (Charlesworth et al., 2018; Gurbich & Bachtrog, 2008; Meisel & Connallon, 2013; Rice, 1984; Vicoso & Charlesworth, 2006).

Since males have only one X chromosome, whereas females have two, X chromosomes differ from autosomes in key population parameters. In a population with equal sex ratio and  $N$  individuals, there are  $1.5N$  X chromosomes and  $2N$  sets of autosomes, so the population size of an X chromosome is three-quarters of the population size of an autosome. Furthermore, X chromosomes are transmitted two-thirds of the time through females and one-third of the time through males, whereas autosomes spend an equal amount of time in males and females. Finally, once Y-linked genes have been lost, recessive mutations arising on an X chromosome are immediately selected in hemizygous males. How these peculiarities affect the evolutionary dynamics of X-linked loci has been previously modeled (Charlesworth et al., 1987; Hitchcock & Gardner, 2020; Meisel & Connallon, 2013; Patten, 2019; Rice, 1984; Vicoso & Charlesworth, 2009). Rice (1984) found that new recessive, male-beneficial sexually antagonistic mutations (i.e., mutations with opposite fitness effects in males and females) can invade a population more easily if they are X-linked than autosomal, and suggested that this may lead to an excess of X-linked genes underlying sexual dimorphism. However, this prediction depends on the dominance coefficient of sexually antagonistic mutations (Rice, 1984), for which we have little empirical evidence, so it is hard to make clear predictions as to whether the X or autosomes are more favorable to the invasion of sexually antagonistic mutations (Ruzicka & Connallon, 2020). Charlesworth et al. (1987) further showed that selection on recessive mutations is stronger on the X chromosome than on the autosomes, resulting in faster substitution rates of recessive beneficial mutations on the X chromosome than on the autosomes, a pattern known as the “faster-X effect” (Vicoso & Charlesworth, 2006, 2009). Conversely, they found that slower substitution rates are expected on X-linked loci for deleterious recessive mutations. Mutations with stronger effects on male than female fitness are particularly prone to faster-X evolution (whereas mutations with female-limited effects are exempt). Extensions of this theory have further shown that male-biased mutation rates (Kirkpatrick & Hall, 2004) and increased variance in male relative to female reproductive success (through its effect on X and autosome effective population size) can increase the faster-X effect (Vicoso & Charlesworth, 2009). On the other hand, if positive selection acts on standing variation rather than new mutations, faster-X evolution is not expected (Orr & Betancourt, 2001). Several studies have attempted to detect a faster-X effect empirically, for instance by testing for a higher proportion of adaptive substitutions on the X chromosome compared to autosomes. Faster-X divergence and faster-X adaptation have been found in various vertebrate and invertebrate clades (Bechsgaard et al., 2019; Charlesworth et al., 2018; Llopart, 2018; Meisel & Connallon, 2013; Mongue et al., 2022; Rupp et al., 2017; Sackton et al., 2014), but not all (Pinharanda et al., 2019; Rousselle et al., 2016; Radhakrishnan & Valenzuela, 2017). Similarly, an excess of sexually antagonistic effects has been suggested in some studies (Abbott et al., 2020; Foerster et al., 2007; Gibson et al., 2002; Innocenti & Morrow, 2010) but not others (Fry, 2010; Ruzicka et al., 2019; Ruzicka & Connallon, 2022). This is further complicated by the fact that the quantitative genetic

measures typically used to detect sexual antagonism are biased toward the detection of X-linked effects (Ruzicka & Connallon, 2020). Because these theories predict different adaptive trajectories for the X and autosomes, they have been invoked to account for various unusual patterns observed on X chromosomes, such as the differential representation of genes with sex-biased expression or the excess movement out of the X (reviewed in Gurbich & Bachtrog, 2008; Vicoso & Charlesworth, 2006).

Much less is known about the evolution of X-linked genes during the early stages of sex chromosome evolution, when the majority of genes on the Y chromosome are functional, but recombination between X and the Y is suppressed. The evolution of X-linked genes under such a scenario (in which there is no recombination between homologous loci on the X and Y, but both X and Y homologs are functional and affect fitness), has not been theoretically explored (but see Engelstaedter, 2008, who modeled how the accumulation of deleterious mutations on the X affects Y-chromosome evolution). By contrast, several empirical studies recognized that young, diploid X-linked loci can have unusual evolutionary dynamics (Nozawa et al., 2016, 2021; Wright et al., 2017). Faster rates of nonsynonymous substitutions and enrichment of sex-biased genes have been detected on young and undifferentiated X/Z chromosomes (Pucholt et al., 2017; Wright et al., 2017). Furthermore, Nozawa et al. (2016, 2021) found evidence of accelerated pseudogenization rates on the young X chromosomes of several *Drosophila* lineages, compared with both the ancient X chromosome and the autosomes. They hypothesized that mechanisms similar to the ones causing degeneration of the Y chromosome could be driving the degeneration of the X chromosome: first, the X chromosome has a smaller population size compared to the autosomes, which makes selection less efficient. Second, since X chromosomes do not recombine in males, their effective population size can be further reduced. However, the opposite is expected in *Drosophila*, as in this clade recombination is restricted to females, where X chromosomes are found two third of the time. Third, female-biased transmission could be driving the loss of genes that are unimportant for females. Another effect that could contribute to the accumulation of deleterious mutations on young X-linked genes is sheltering by the functional gene copy on the Y chromosome. New mutations arising on a diploid X-locus are always heterozygous in males because there is no recombination between the X and the Y, and their phenotypic effect is masked by the ancestral allele on the Y. While the role of sheltering has been appreciated in other contexts, such as the degeneration of Y chromosomes (Bachtrog, 2013; Muller, 1914; Nei, 1970) and the evolution of recombination suppression (Antonovics & Abrams, 2004; Charlesworth & Wall, 1999; Jay et al., 2022; Olito et al., 2022), it is unclear to what extent it affects early X-chromosome evolution. Here, we model evolutionary rates of X-linked loci with functional Y copies under various selective regimes, to formally explore how these different processes shape the early stages of X chromosome evolution.

## Methods

### The diffusion approximation

Substitution rates can be calculated as the average number of mutations entering a population in one generation times the fixation probability of those mutations (Kimura & Ohta, 1971). To derive probabilities of fixation for autosomal and hemizygous X-linked loci, Vicoso and Charlesworth (2009) used the diffusion approximation. Here, we extend their model to diploid X-linked loci.

Let  $A_1$  and  $A_2$  be alleles for some locus, with frequencies  $(1-p)$  and  $p$ , respectively, and fitness effects as noted in Table 1.

We can estimate the fixation probability of allele  $A_2$  using the diffusion approximation (Ewens, 2004). The fixation probability of an allele with the initial frequency  $p$  is given by the function  $U(p)$ :

$$U(p) = \frac{\int_0^p G(y)dy}{\int_0^1 G(y)dy}, \tag{1}$$

With  $G(y) = \exp(-2 \int_0^y \frac{M(x)}{V(x)} dx)$ ,

where  $M(x)$  and  $V(x)$  are, respectively, the expectation and the variance of the change of allele frequency.

Assuming a weak effect of selection in each sex, so that second-order terms are small enough to be neglected, the fitness of a genotype can be approximated as the average of fitness effects in males and females (Charlesworth & Charlesworth, 2010, ch. 3.1). For a diploid X-linked locus, the expected change of an allele  $A_2$  frequency due to selection is given by:

$$M_{dx}(x) \approx x(1-x) \left( \frac{2}{3}(w_{2f} - w_{1f}) + \frac{1}{3}(w_{2m} - w_{1m}) \right),$$

where  $x$  is the frequency of the allele  $A_2$ ,  $w_{1f}$  and  $w_{1m}$  are marginal fitnesses of allele  $A_1$  in females and males, respectively, and  $w_{2f}$  and  $w_{2m}$  are marginal fitnesses of allele  $A_2$  in females and males respectively. Calculating these marginal fitnesses from Table 1, we have:

$$M_{dx}(x) \approx x(1-x) \left( \frac{2}{3}s_f(h + x(1-2h)) + \frac{1}{3}hs_m \right),$$

where  $h$  is the dominance coefficient.

We now divide  $M_{dx}(x)$  by the variance in the change of allele  $A_2$  frequency due to sampling drift,

$$V_{dx}(x) \approx \frac{x(1-x)}{2N_{ex}},$$

where  $N_{ex}$  is the effective population size of the X chromosome, and integrate to find  $G(y)$ . This can be written in terms of mean selection averaged across sexes,

$\bar{s} = \frac{2}{3}s_f + \frac{1}{3}hs_m = \frac{s_f + hs_m}{3}$ , and  $\sigma = \frac{s_f(1-2h)}{3}$ , which is zero if there is no dominance ( $h = 1/2$ ). Then:

$$\begin{aligned} G_{dx}(y) &= \exp\left(-2 \int_0^y \frac{M(x)}{V(x)} dx\right) \\ &= \exp\left(-\frac{4}{3}N_{ex}(2s_f + s_m)h \int_0^y dx - \frac{8}{3}N_{ex}s_f(1-2h) \int_0^y x dx\right) \\ &= \exp\left(-\frac{4}{3}N_{ex}(2s_f + s_m)hy - \frac{4}{3}N_{ex}s_f(1-2h)y^2\right) \\ &= \exp\left(-4N_{ex}\left(\frac{1}{3}(s_f + hs_m)y - \frac{1}{3}s_f(1-2h)y(1-y)\right)\right) \\ &= \exp\left(-4N_{ex}(\bar{s}y - \sigma y(1-y))\right) \end{aligned} \tag{2}$$

We are interested in the probability of fixation of a single new mutation, which is initially at  $p = \frac{2}{3N}$ ; the substitution rate is

$1.5N\mu U(p) = \mu U(p)/p$ , where  $\mu$  is the mutation rate. Following Eq. (1), the fixation probability  $U(p)$  becomes:

$$\begin{aligned} U(p) &= \frac{\int_0^p G(y)dy}{\int_0^1 G(y)dy} \\ &= \frac{\operatorname{erf}\left(\frac{N_{ex}(\bar{s} + \sigma(2p-1))}{\sqrt{N_{ex}\sigma}}\right) - \operatorname{erf}\left(\frac{N_{ex}(\bar{s} - \sigma)}{\sqrt{N_{ex}\sigma}}\right)}{\operatorname{erf}\left(\frac{N_{ex}(\bar{s} - \sigma)}{\sqrt{N_{ex}\sigma}}\right) - \operatorname{erf}\left(\frac{N_{ex}(\bar{s} - \sigma)}{\sqrt{N_{ex}\sigma}}\right)} \\ &= \frac{\frac{4p\sqrt{N_{ex}\sigma}}{\sqrt{\pi}} \exp\left(-\frac{N_{ex}(\bar{s} - \sigma)^2}{\sigma}\right)}{\operatorname{erf}\left(\frac{N_{ex}(\bar{s} - \sigma)}{\sqrt{N_{ex}\sigma}}\right) - \operatorname{erf}\left(\frac{N_{ex}(\bar{s} - \sigma)}{\sqrt{N_{ex}\sigma}}\right)} + O(p^2), \end{aligned}$$

where  $O(p^2)$  are higher order terms in the Taylor series when  $p$  is close to zero. Assuming weak selection and sufficiently large effective population size,  $O(p^2)$  can be neglected for new mutations. Then, the first-order approximation in  $p$  for the substitution rate is:

$$\mu(U(p)/p) = 4\mu \frac{\sqrt{N_{ex}\sigma}}{\pi} \frac{\exp(-N_{ex}(\bar{s} - \sigma)^2/\sigma)}{(\operatorname{erf}(N_{ex}(\bar{s} + \sigma)/\sqrt{N_{ex}\sigma}) - \operatorname{erf}(N_{ex}(\bar{s} - \sigma)/\sqrt{N_{ex}\sigma}))} \tag{3}$$

Note that the substitution rate relative to mutation depends only on  $N_{ex}\bar{s}$  and  $N_{ex}\sigma$ ; in the additive case ( $h = 1/2$ ), it simplifies to Kimura's formula,  $4N_{ex}\bar{s}/(1 - \exp(-4N_{ex}\bar{s}))$ .

Also, note from Table 1 that the only difference between the hemizygous and the diploid X cases is that  $s_m$  is replaced by  $hs_m$  in the latter case; thus, the results of Vicoso and Charlesworth (2009) for the hemizygous case can be found from the results given here simply by replacing  $s_m$  by  $s_m/h$ .


We implemented the numerical integration in R (version 4.1.1, code available at [https://github.com/andrea-mrnjavac/X-chromosome-theory/blob/main/fixation\\_probability\\_functions.R](https://github.com/andrea-mrnjavac/X-chromosome-theory/blob/main/fixation_probability_functions.R)), which allowed us to derive substitution rates for autosomal, hemizygous X-linked and diploid X-linked loci over the range of dominance coefficients and selective effects in males and females. A GUI application with implemented fixation probability functions for autosomal, hemizygous X-linked, and diploid X-linked loci is available at <https://degenerate-x.science.ista.ac.at/>, and allows the user to explore the difference among substitution rates of autosomal, hemizygous X-linked and diploid X-linked loci over the range of parameter values (selective effects in males and females and dominance coefficient), assuming effective population size of an X chromosome is three fourth of autosomal effective population size. In addition, we modeled evolutionary rates on X chromosomes and autosomes assuming equal population sizes for autosomes and X chromosomes, in order to disentangle the effects of effective population size and sheltering.

### Branching process approximation

To model the substitution rates of strongly beneficial mutations at diploid X-linked loci with functional but non-recombining Y

**Table 1.** Relative fitnesses in females and males for autosomal, hemizygous X-linked, and diploid X-linked loci.

		Females			Males		
Autosomal							
	Genotypes	$A_1A_1$	$A_1A_2$	$A_2A_2$	$A_1A_1$	$A_1A_2$	$A_2A_2$
	Fitness	1	$1+hs_f$	$1+s_f$	1	$1+hs_m$	$1+s_m$
Hemizygous X-linked							
	Genotypes	$A_{x1}A_{x1}$	$A_{x1}A_{x2}$	$A_{x2}A_{x2}$	$A_{x1}$	$A_{x2}$	
	Fitness	1	$1+hs_f$	$1+s_f$	1	$1+s_m$	
Diploid X-linked							
	Genotypes	$A_{x1}A_{x1}$	$A_{x1}A_{x2}$	$A_{x2}A_{x2}$	$A_{y1}A_{x1}$	$A_{y1}A_{x2}$	
	Fitness	1	$1+hs_f$	$1+s_f$	1	$1+hs_m$	



**Figure 1.** Ratios of substitution rates ( $R$ ) on hemizygous X to autosomal loci and diploid X to autosomal loci as functions of dominance coefficient  $h$ , for beneficial,  $N_{eA}s = 3$  (A) and deleterious,  $N_{eA}s = -1$  mutations (B).

gametolog and allow comparisons with the classic Charlesworth et al. (1987) result, we used Haldane’s branching process approximation (Charlesworth et al., 1987, 2018; Haldane, 1927; Meisel & Connallon, 2013; Vicoso & Charlesworth, 2006).

If  $N_e sh$  is sufficiently large ( $1/N_e \ll sh \ll 1$ ), the fixation probability of a new beneficial mutation can be approximated as twice the advantage of a heterozygous genotype (Haldane, 1927).

Taking into account the effects of a mutation in males and females separately (Vicoso & Charlesworth, 2006), the fixation probability of a single new beneficial mutation at an autosomal locus can be approximated as:

$$P_A\left(\frac{1}{2N}\right) \approx \frac{1}{2}2hs_f + \frac{1}{2}2hs_m \approx h(s_f + s_m),$$

where  $s_f$  and  $s_m$  are selection coefficients in females and males, respectively,  $h$  is the dominance coefficient, and  $N$  is the number of diploid individuals, as before.

For diploid X-linked loci, the fixation probability of a new mutation is then:

$$P_X\left(\frac{1}{1.5N}\right) \approx \frac{2}{3}2hs_f + \frac{1}{3}2hs_m \approx \frac{2}{3}h(2s_f + s_m).$$

If we assume the number of X chromosomes in a population is three fourth the number of autosomes, and dominance is the same in males and females, as well as the mutation rate, the following substitution rates are then:

$$K_A \approx 2N\mu h(s_f + s_m) \text{ for autosomal loci, and}$$

$$K_X \approx \frac{3}{2}N\mu \frac{2}{3}h(2s_f + s_m) \approx N\mu h(2s_f + s_m) \text{ for diploid X loci.}$$

The ratio of diploid X to autosomal adaptive substitution rates is:

$$R \approx \frac{N\mu h(2s_f + s_m)}{2N\mu h(s_f + s_m)} \approx \frac{2s_f + s_m}{2s_f + 2s_m} \quad (4)$$

## Results

### Diploid X loci adapt slower and accumulate deleterious mutations faster


In addition to previously derived substitution rates on hemizygous X loci and autosomal loci (Charlesworth et al., 1987; Vicoso & Charlesworth, 2009), we used the diffusion approximation to estimate fixation probabilities of

new mutations arising at diploid X-linked loci (which have a functional homolog on the Y), and their substitution rates.

We verified these estimates with individual-based forward genetic simulations in SLiM (Haller & Messer, 2019) (Supplementary Figure 1, details of simulations are given in Supplementary Material). The X/A ratios of substitution rates ( $R$ ) for beneficial and deleterious mutations and for diploid and hemizygous X-linked loci are visualized in Figure 1. We recover the previously described faster-X effect for hemizygous X-linked loci (Charlesworth et al., 1987; Vicoso & Charlesworth, 2006, 2009), where recessive beneficial mutations accumulate faster on the hemizygous X loci, while dominant beneficial mutations accumulate faster on the autosomes compared to the hemizygous X loci. On the contrary, diploid X loci exhibit a slower-X effect regardless of the dominance coefficient: the substitution rate of beneficial mutations is lower at diploid X loci compared to autosomal and hemizygous X loci, in agreement with reduced efficiency of selection on diploid X loci due to sheltering and reduced effective population size (Table 1). Substitution rates of deleterious mutations follow the opposite pattern, with diploid X loci accumulating deleterious mutations faster than autosomal and hemizygous X loci. These results, therefore, show that young diploid X-linked genes have reduced adaptive potential, and increased accumulation of deleterious mutations, relative to both hemizygous X-linked genes and autosomal genes.

To disentangle the effects of reduced effective population size and sheltering on substitution rates of diploid X loci, we calculated the ratio of substitution rates in the case where  $N_{eX} = N_{eA}$ , as is approximately the case in *Drosophila melanogaster* due to the effect of selection at linked sites (although differences in male and female fitness variance may also play a role (Charlesworth, 2001)): since there is no recombination in males, X chromosomes have higher effective recombination rates than autosomes, which increases the  $N_{eX}/N_{eA}$  ratio (Charlesworth et al., 2018). The results show that in this case there is still a “slower-X” effect for diploid X loci, but only for recessive mutations (Supplementary Figure 4).

As an example, Figure 1 shows the ratios of substitution rates ( $R$ ) for beneficial mutations with  $N_{eA}s = 3$ , and for deleterious mutations with  $N_{eA}s = -1$ , where equal effects in males and females are assumed ( $s = s_f = s_m$ ), but a wide range of positive and negative  $N_{eA}s$  values (we modeled  $N_{eA}s$  values from  $-3$  to  $5$ ) yield



**Figure 2.** Ratios of X/A substitution rates ( $R$ ) of male-limited mutations ( $N_{e_s m} = 3, N_{e_s f} = 0$  for and  $N_{e_s m} = -1, N_{e_s f} = 0$  for beneficial and deleterious mutations respectively) are plotted alongside ratios of substitution rates of mutations with equal effects ( $N_{e_s m} = 3, N_{e_s f} = 3$  for and  $N_{e_s m} = -1, N_{e_s f} = -1$  for beneficial and deleterious mutations respectively), and female-limited ( $N_{e_s m} = 0, N_{e_s f} = 3$  for and  $N_{e_s m} = 0, N_{e_s f} = -1$  for beneficial and deleterious mutations, respectively) effects for beneficial (A) and deleterious (B) mutations, as a function of dominance coefficient,  $h$ .  $R$  for female-biased mutations is between  $R$  for female-limited mutations and  $R$  for mutations with equal effects in males and females, while  $R$  for male-biased mutations is between  $R$  for male-limited mutations and  $R$  for mutations with equal effects in both sexes.

the same qualitative pattern. The effect of different  $N_{e_s}$  values of a mutation on the X/A ratio of substitution rates can be explored in a GUI web application provided at <https://degenerate-x.science.ista.ac.at/>. However, it is worth noting that the ratio of substitution rates at diploid X loci compared to autosomal loci ( $R$ ) increases exponentially with the strength of the deleterious effect of a mutation (Supplementary Figure 2). This means that mutations with stronger deleterious effects will be fixed much more often in diploid X loci relative to an autosome.

While the original faster-X publication (Charlesworth et al., 1987) focused on  $R$  (in that case the autosome:X substitution rate ratio), evolutionary rates are usually measured as the rate of nonsynonymous substitutions ( $dN$ ), normalized by the synonymous rate of substitutions ( $dN/dS$ , with  $dS$  acting as a proxy for neutral divergence) (Meisel & Connallon, 2013). To facilitate comparisons with such data, we also plot the substitution rate of beneficial and deleterious mutations after normalizing them by the neutral substitution rate (qualitative patterns remain the same, Supplementary Figure 3).

Similar results are recovered with the branching process approximation (see methods). Equation (4) shows that for strongly beneficial mutations (where  $N_e s h$  is sufficiently large), the X/A ratio of substitution rates does not depend on the dominance coefficient at diploid X loci, in contrast with the previously described X/A ratio of adaptive evolutionary rates at hemizygous X loci, which exhibit faster-X effect for recessive mutations. More precisely, our results show that for mutations with female-limited selective effects, substitution rates are the same at autosomal and diploid X loci, that is,  $R = 1$ ; for mutations with male-limited selective effects,  $R = 1/2$ , that is, adaptation rate on diploid X is half the adaptation rate on autosomes, and for mutations with equal effects in males and females,  $R = 3/4$ . These results indicate a slower-X effect for all strongly beneficial mutations arising on diploid X loci and having an effect in males.

### The “slower X” effect is strongest for male-biased mutations

We also aimed to disentangle how selective effects in males and females separately affect the substitution rate at diploid X loci.

Mutations can have different effects in males and females: they can be sex-limited, affecting only the fitness of one sex, sex-biased, if they have a stronger effect on the fitness of one sex than the other, or sexually antagonistic, if they have fitness effects of opposite signs in the two sexes. We can intuitively see from the fitness table (Table 1) that the differences in evolutionary rates of diploid X, hemizygous X, and autosomal loci result from the differences in the male part of the fitness table. Indeed, similar to Charlesworth et al. (1987), we find that evolutionary rates are the same at autosomal, hemizygous X-linked, and diploid X-linked loci for female-limited mutations (mutations affecting only female fitness, e.g.,  $N_{e_s m} = 0, N_{e_s f} = 3$  in Figure 2A, or  $N_{e_s m} = 0, N_{e_s f} = -1$  in Figure 2B). On the other hand, the X/A ratio of substitution rates ( $R$ ) for beneficial male-limited mutations ( $N_{e_s m} = 3, N_{e_s f} = 0$ ), and to a smaller extent male-biased mutations, is lower than  $R$  for mutations with an equal effect in both males and females ( $N_{e_s m} = 3, N_{e_s f} = 3$ ) (Figure 2A). Male-limited and male-biased mutations are primarily under selection in males, where their effect is masked by the ancestral allele on the Y, resulting in a stronger “slower-X effect.” Analogously, deleterious male-limited and male-biased mutations accumulate faster at diploid X loci than at autosomal loci, and faster than deleterious female-limited and female-biased loci with the corresponding fitness effects (e.g., comparing male-biased mutations with  $N_{e_s m} = -1, N_{e_s f} = -0.5$  to female-biased mutations with  $N_{e_s m} = -0.5, N_{e_s f} = -1$ ). Counterintuitively, Figure 2B shows that  $R$  for a diploid X is larger for mutations with equal fitness effects in males and females than for male-limited and male-biased mutations. This is because mutations affecting both sexes ( $N_{e_s m} = -1, N_{e_s f} = -1$ ) are overall more deleterious than mutations affecting males ( $N_{e_s m} = -1, N_{e_s f} = 0$ ) or females only, and  $R$  for the diploid X increases exponentially with the strength of the deleterious effect of a mutation (Supplementary Figure 2). To summarize, the X/A ratio of substitution rates for deleterious mutations at diploid X will be greater than 1 as long as the mutation has an effect in males, and mutations with a stronger deleterious effect in males than in females will accumulate in excess on the young X compared to autosomes.

Sexually antagonistic mutations are expected to accumulate faster on an ancient hemizygous X than on the autosomes if they are recessive and male-beneficial, or dominant and female-beneficial (Rice, 1984; Vicoso & Charlesworth, 2009, and Figure 3). One assumption of these studies, which we also make here, is that the dominance coefficient of antagonistic mutations is the same in males and females (see Fry, 2010, for results when this does not hold). The resulting differential accumulation of sexually antagonistic mutations has been proposed to influence the evolution of sex chromosomes and their role in encoding sexual dimorphism (Fry, 2010; Rice, 1984). Figure 3 shows that mutations providing an advantage to males but a disadvantage to females ( $N_{eS_m} = 3$ ,  $N_{eS_f} = -3$ ) accumulate much more slowly at diploid X loci than at autosomal or hemizygous X loci, independent of their dominance coefficient. Mutations carrying female advantage and male disadvantage ( $N_{eS_m} = -3$ ,  $N_{eS_f} = 3$ ), on the other hand, accumulate faster at diploid X loci. Taken together, these results show that the reduced efficacy of selection on a diploid X is highly influenced by sex-specific fitness effects, with mutations that benefit primarily males tending to accumulate slower, and mutations that are detrimental to males accumulating faster whether they benefit females or not.


## Discussion

There has been extensive theoretical and empirical work on the “faster-X effect” expected on X chromosomes with a degenerated Y counterpart (reviewed in Charlesworth et al., 2018; Meisel & Connallon, 2013). Here, we modeled evolutionary rates of diploid X-linked loci, with a functional, but non-recombining, gametolog on the Y counterpart. Our results show very different evolutionary dynamics for diploid and hemizygous X loci. We make two key predictions for the evolution of young X-chromosomes: (a) Selection efficacy is reduced, such that fewer beneficial mutations and more deleterious mutations will fix there compared to an autosomal locus. (b) The relaxation of selection is stronger for mutations that primarily affect male fitness and/or benefit males at the expense of females. Over time, this may lead to the “demasculinization” of the young X

chromosome, i.e., the degeneration of genes with male-important functions, and/or the failure to acquire new genes with male functions. This is in contrast to hemizygous X-linked loci on differentiated X chromosomes, which may exhibit faster adaptation and masculinization (as long as beneficial mutations are generally recessive).

These peculiar evolutionary dynamics of diploid X loci are caused by: (a) a smaller effective population size compared to autosomes, (b) female-biased transmission, and (c) sheltering of partly recessive X-linked mutations in males by an ancestral allele on the Y. By exploring a range of parameters, we could to some extent quantify the individual contribution of these effects. Sheltering does not affect fully dominant mutations, and the difference in  $R$  when  $h = 0$  and  $h = 1$  shows that it can have a substantial effect on rates of adaptive and maladaptive divergence (Figures 1–3). When the reduction in  $N_e$  is removed ( $N_{eA} = N_{eX}$ ), reduced efficacy of selection is only detected when mutations are at least partly recessive (due to sheltering, Supplementary Figure 4A and B) and/or have male-biased or male-limited effects (due to female-biased transmission and a stronger effect of sheltering, Supplementary Figure 4C and D). The relative importance of these effects depends on the strength of fitness effects in males, as sheltering only affects mutations expressed in this sex. Finally, both female-biased transmission and sheltering contribute to the proposed demasculinization, as male-important mutations are affected disproportionately (Supplementary Figure 4).

Only a few empirical studies have explicitly compared the evolution of diploid X-linked genes to autosomal control. Several of them investigated the young neo-X chromosome found in the *Drosophila miranda* lineage, where a pair of autosomes fused with the ancestral Y chromosome and became a neo-X and neo-Y a little over 1 million years ago. Around 40% of genes on the neo-Y are still functional (Nozawa et al., 2016; Zhou & Bachtrog, 2012). Zhou and Bachtrog (2012) found that hemizygous X loci adapt faster than diploid X loci on the neo-X chromosome in *D. miranda*. Nozawa et al. (2016) further detected evidence of an accelerated pseudogenization rate on the neo-X chromosome in *D. miranda* after it became X-linked. They compared the neo-X chromosome



**Figure 3.** Ratios of X/A substitution rates ( $R$ ) for sexually antagonistic mutations carrying male advantage and female disadvantage ( $N_{eS_m} = 3$ ,  $N_{eS_f} = -3$ ) (A) and mutations carrying female advantage and male disadvantage  $N_{eS_m} = -3$ ,  $N_{eS_f} = 3$  (B) plotted along the  $R$  for mutations with equal effects in males and females, as a function of dominance coefficient,  $h$ .

in *D. miranda* to the corresponding autosome in *D. pseudoobscura*, and found that genes that were ancestrally under reduced selective constraints (have higher  $dN/dS$  values in *D. pseudoobscura*) and genes with ancestrally male-biased function (approximated from *D. pseudoobscura* F/M expression ratios) were more likely to be pseudogenized on the neo-X than we would expect if it was an autosome. Their analyses suggest that the reduction in efficiency of selection, especially for male-biased genes, is causing the accelerated pseudogenization rate on the young X chromosome. Recently, Nozawa et al. (2021) found degeneration of neo-X chromosomes in two other *Drosophila* species with independently acquired neo-X. These results are generally in line with our theoretical predictions of the maladaptive evolution of young X chromosomes and accompanying demasculinization, but work in various other systems is needed to understand how universal this pattern is and how much of a contribution it makes to X-chromosome evolution.

In particular, one question that we did not address here is the time frame over which maladaptive evolution occurs. In the early stages of sex chromosome differentiation, the majority of the ancestral gene content on the Y chromosome will still be functional, and corresponding X-linked loci will be diploid and evolve under reduced selective efficacy. The accelerated rate of pseudogenization on the X will slow down as the Y chromosome degenerates and more X-linked loci become hemizygous, causing a shift in evolutionary dynamics to standard “faster-X.” Theory predicts that Y chromosomes first degenerate quickly after the recombination suppression but after they have lost about half of the gene content the process slows down, since the rates of degeneration by Muller’s ratchet, background selection, and genetic hitchhiking correlate with the number of active genes (Bachtrog, 2008). For instance, more than half of the genes on the *Drosophila miranda* neo-Y chromosome have been lost in a little over 1 million years, and the 15-million-year-old neo-Y of *D. pseudoobscura* is already highly degenerated (reviewed in Charlesworth, 2021). However, it is unclear if the neo-sex chromosomes of *Drosophila*, which have quickly co-opted a preexisting mechanism of dosage compensation, are representative of typical dynamics of Y degeneration, and much slower Y degeneration has been described in other systems (Charlesworth, 2021; Li et al., 2021). Nonrecombining regions with intermediate or low levels of Y/W degeneration have been described in various taxa (Charlesworth, 2021), e.g.: schistosomes (Elkrewi et al., 2021), frogs (Furman & Evans, 2018), crustaceans (Elkrewi et al., 2022), birds (Liu et al., 2021), fish (Sardell et al., 2021), and plants (Veltos et al., 2019). It is therefore clear that many young X-linked genes can remain diploid for substantial periods of time.

The existence of a period of maladaptive evolution has implications for young homomorphic X chromosomes, but may also contribute to patterns observed on older sex chromosomes. The fact that X-linked genes with male-specific functions should be more prone to early maladaptive evolution suggests that the “demasculinization” that is observed on differentiated X-chromosomes of various species (Gurbich & Bachtrog, 2008) may begin before the degeneration of the Y. The subsequent shift in evolutionary dynamics could contribute to the differences in sex-biased gene contents of X chromosomes, which are for instance masculinized in mammals and demasculinized in *Drosophila* lineage (Gurbich & Bachtrog, 2008). The importance of temporal dynamics to the process of demasculinization is well appreciated in *Drosophila*, where only ancient male-biased genes are depleted from the X chromosome, whereas newly evolved ones are enriched on it (Zhang et al., 2010). While this potentially supports a role of

diploid X evolution, the explanations brought forward to explain it typically assume a hemizygous X. Furthermore, Engelstaeder (2008) used modeling to show that the accumulation of deleterious mutations on diploid X-linked loci can slow down the degeneration of their Y-linked loci. An illustration of this is PRSSLY, a gene that was lost from the mammalian X chromosome in eutherians but retained on the Y (Hughes et al., 2022). If genes that function primarily in males are the ones that tend to accumulate deleterious mutations on a young X, this may lead to their preferential maintenance on degenerating Y chromosomes. The preservation of genes with male-biased expression on the Y has been observed (Crowson et al., 2017; Kaiser et al., 2011; Mahajan & Bachtrog, 2017; Zhou & Bachtrog, 2012) and is usually assumed to be driven by male-specific selection on Y-linked genes. In fact, degeneration of male-important genes on the X may drive their conservation on the Y, as well as the other way around.

Other peculiarities of X chromosomes may first arise early in their evolution. Repetitive sequences and transposable elements are overrepresented on X chromosomes as well as Y chromosomes (Bellott et al., 2010). The reduction in the effective population size of young X chromosomes (due to their lower population size, and potentially further exacerbated because they do not recombine in males) may already contribute to the accumulation of repeats. Finally, X-chromosomes are central to the two “rules of speciation”: Haldane’s rule (hybrid sterility or inviability tends to affect the heterogametic sex more than the homogametic sex), and the “large-X effect” (an excessive proportion of hybrid sterility loci maps to X chromosomes). Most of the clades that obey these rules have differentiated sex chromosomes, and explanations have typically invoked the faster-X hypothesis (along with other models) (Dufresnes & Crochet, 2022). However, hybridization patterns in *Aedes* mosquitoes, which have undifferentiated sex chromosomes, follow Haldane’s rule (Presgraves & Orr, 1998). Similarly, Dufresne et al. (2016) detected an excessive role of an undifferentiated X-chromosome in the reproductive isolation of tree frogs, and these rules have been suggested to apply to some clades with homomorphic sex chromosomes (Filatov, 2018). An excessive accumulation of deleterious mutations on homomorphic X chromosomes may provide an explanation for a large-X effect in this context, if compensatory mutations arise elsewhere in the genome. Similarly, if Y-degradation and X-sheltering affect different genes in close species, a mismatch between them may contribute to male sterility in hybrids. Studies of hybrids over a wide range of sex chromosome differentiation will in the future allow us to quantify the temporal dynamics of the large-X effect and Haldane’s rule, and the contribution of diploid X evolution.

In short, X-linked loci are expected to undergo a period of maladaptive evolution and demasculinization in the early stages of their differentiation. Our results show that contrary to what is often assumed, the peculiar evolutionary patterns on the X chromosome may arise before substantial degeneration of the Y has occurred and provide a novel framework for interpreting the increasing amount of data available for clades with young sex chromosomes.

## Supplementary material

Supplementary material is available online at *Evolution Letters* (<https://academic.oup.com/evlett/qrac004>)

## Data availability

No data was created in this project.

## Author contributions

A.M. and B.V. designed the study and wrote the first draft. A.M., K.K., and N.B. performed the analysis. All the authors contributed to and approved the final version of the manuscript.

## Conflict of interest

The authors declare no conflict of interest.

## Acknowledgments

We thank the Vicoso and Barton groups and ISTA Scientific Computing Unit. We also thank two anonymous reviewers for their valuable comments. This work was supported by the European Research Council under the European Union's Horizon 2020 research and innovation program (grant agreements no. 715257 and no. 716117).

## References

- Abbott, J. K., Chippindale, A. K., & Morrow, E. H. (2020). The microevolutionary response to male-limited X-chromosome evolution in *Drosophila melanogaster* reflects macroevolutionary patterns. *Journal of Evolutionary Biology*, **33**, 738–750.
- Antonovics, J., & Abrams, J. Y. (2004). Intratetrad mating and the evolution of linkage relationships. *Evolution*, **58**, 702–709.
- Bachtrog, D. (2008). The temporal dynamics of processes underlying Y chromosome degeneration. *Genetics*, **179**, 1513–1525.
- Bachtrog, D. (2013). Y-chromosome evolution: Emerging insights into processes of Y-chromosome degeneration. *Nature Reviews. Genetics*, **14**, 113–124.
- Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T.-L., Hahn, M. W., Kitano, J., Mayrose, I., Ming, R., Perrin, N., Ross, L., Valenzuela, N., Vamosi, J. C. & The Tree of Sex Consortium. (2014) Sex determination: Why so many ways of doing it? *PLOS Biology*, **12**, e1001899.
- Bechsgaard, J., Schou, M. F., Vanthournout, B., Hendrickx, F., Knudsen, B., Settepani, V., Schierup, M. H. & Bilde, T. (2019) Evidence for faster X chromosome evolution in spiders. *Molecular Biology and Evolution*, **36**, 1281–1293.
- Bellott, D. W., Skaletsky, H., Pyntikova, T., Mardis, E. R., Graves, T., Kremitzki, C., Brown, L. G., Rozen, S., Warren, W. C., Wilson, R. K. & Page, D. C. (2010) Convergent evolution of chicken Z and human X chromosomes by expansion and gene acquisition. *Nature*, **466**, 612–616.
- Charlesworth, B. (2001). The effect of life-history and mode of inheritance on neutral genetic variability. *Genetics Research*, **77**, 153–166.
- Charlesworth, D. (2021). The timing of genetic degeneration of sex chromosomes. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **376**, 20200093.
- Charlesworth, B., Campos, J. L., & Jackson, B. C. (2018). Faster-X evolution: Theory and evidence from *Drosophila*. *Molecular Ecology*, **27**, 3753–3771.
- Charlesworth, B., & Charlesworth, D. (2010). *Elements of evolutionary genetics*. W. H. Freeman.
- Charlesworth, B., Coyne, J. A., & Barton, N. H. (1987). The relative rates of evolution of sex chromosomes and autosomes. *The American Naturalist*, **130**, 113–146.
- Charlesworth, B., & Wall, J. D. (1999). Inbreeding, heterozygote advantage and the evolution of neo-X and neo-Y sex chromosomes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **266**, 51–56.
- Crowson, D., Barrett, S. C. H., & Wright, S. I. (2017). Purifying and positive selection influence patterns of gene loss and gene expression in the evolution of a plant sex chromosome system. *Molecular Biology and Evolution*, **34**, 1140–1154.
- Dufresnes, C., & Crochet, P.-A. (2022). Sex chromosomes as supergenes of speciation: Why amphibians defy the rules? *Philosophical Transactions of the Royal Society B: Biological Sciences*, **377**, 20210202.
- Dufresnes, C., Majtyka, T., Baird, S. J. E., Gerchen, J. F., Borzée, A., Savary, R., Ogielska, M., Perrin, N. & Stöck, M. (2016). Empirical evidence for large X-effects in animals with undifferentiated sex chromosomes. *Scientific Reports*, **6**, 21029.
- Elkrewi, M., Khauratovich, U., Troups, M. A., Bett, V. K., Mrnjavac, A., Macon, A., Fraisse, C., Sax, L., Huylmans, A. K., Hontoria, F. & Vicoso, B. (2022). ZW sex-chromosome evolution and contagious parthenogenesis in *Artemia* brine shrimp. *Genetics*, **222**, iyac123.
- Elkrewi, M., Moldovan, M. A., Picard, M. A. L., & Vicoso, B. (2021). Schistosome W-linked genes inform temporal dynamics of sex chromosome evolution and suggest candidate for sex determination. *Molecular Biology and Evolution*, **38**, 5345–5358.
- Engelstädter, J. (2008). Muller's ratchet and the degeneration of Y chromosomes: A simulation study. *Genetics*, **180**, 957–967.
- Ewens, W. J. (2004). *Mathematical population genetics. Interdisciplinary applied mathematics*. Springer.
- Filatov, D. A. (2018). The two “rules of speciation” in species with young sex chromosomes. *Molecular Ecology*, **27**, 3799–3810.
- Foerster, K., Coulson, T., Sheldon, B. C., Pemberton, J. M., Clutton-Brock, T. H., & Kruuk, L. E. B. (2007). Sexually antagonistic genetic variation for fitness in red deer. *Nature*, **447**, 1107–1110.
- Fry, J. D. (2010). The genomic location of sexually antagonistic variation: Some cautionary comments. *Evolution*, **64**, 1510–1516.
- Furman, B. L. S., & Evans, B. J. (2018). Divergent evolutionary trajectories of two young, homomorphic, and closely related sex chromosome systems. *Genome Biology and Evolution*, **10**, 742–755.
- Gibson, J. R., Chippindale, A. K., & Rice, W. R. (2002). The X chromosome is a hot spot for sexually antagonistic fitness variation. *Proceedings of the Royal Society B: Biological Sciences*, **269**, 499–505.
- Gurbich, T. A., & Bachtrog, D. (2008). Gene content evolution on the X chromosome. *Current Opinion in Genetics & Development*, **18**, 493–498.
- Haldane, J. B. S. (1927). A mathematical theory of natural and artificial selection, part V: Selection and mutation. *Mathematical Proceedings of the Cambridge Philosophical Society*, **23**, 838–844.
- Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward genetic simulations beyond the Wright–Fisher model. *Molecular Biology and Evolution*, **36**, 632–637.
- Hitchcock, T. J., & Gardner, A. (2020). A gene's-eye view of sexual antagonism. *Proceedings of the Royal Society B: Biological Sciences*, **287**, 20201633.
- Hughes, J. F., Skaletsky, H., Nicholls, P. K., Drake, A., Pyntikova, T., Cho, T.-J., Bellott, D. W. & Page, D. C. (2022). A gene deriving from the ancestral sex chromosomes was lost from the X and retained on the Y chromosome in eutherian mammals. *BMC Biology*, **20**, 133.
- Innocenti, P., & Morrow, E. H. (2010). The sexually antagonistic genes of *Drosophila melanogaster*. *PLOS Biology*, **8**, e1000335.
- Jay, P., Tezenas, E., Véber, A., & Giraud, T. (2022). Sheltering of deleterious mutations explains the stepwise extension of recombination suppression on sex chromosomes and other supergenes. *PLOS Biology*, **20**, e3001698.
- Jeffries, D. L., Gerchen, J. F., Scharmann, M., & Pannell, J. R. (2021). A neutral model for the loss of recombination on sex chromosomes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **376**, 20200096.
- Kaiser, V. B., Zhou, Q., & Bachtrog, D. (2011). Nonrandom gene loss from the *Drosophila miranda* neo-Y chromosome. *Genome Biology and Evolution*, **3**, 1329–1337.



- Kimura, M., & Ohta, T. (1971). On the rate of molecular evolution. *Journal of Molecular Evolution*, **1**, 1–17.
- Kirkpatrick, M., & Hall, D. W. (2004). Male-biased mutation, sex linkage, and the rate of adaptive evolution. *Evolution*, **58**, 437–440.
- Larson, E. L., Kopania, E. E. K., & Good, J. M. (2018). Spermatogenesis and the evolution of mammalian sex chromosomes. *Trends in Genetics*, **34**, 722–732.
- Li, M., Zhang, R., Fan, G., Xu, W., Zhou, Q., Wang, L., Li, W., Pang, Z., Yu, M., Liu, Q., Liu, X., Schartl, M. & Chen, S. (2021). Reconstruction of the origin of a Neo-Y Sex chromosome and its evolution in the spotted knifejaw, *oplegnathus punctatus*. *Molecular Biology and Evolution*, **38**, 2615–2626.
- Liu, J., Wang, Z., Li, J., Xu, L., Liu, J., Feng, S., Guo, C., Chen, S., Ren, Z., Rao, J., Wei, K., Chen, Y., Jarvis, E. D., Zhang, G. & Zhou, Q. (2021). A new emu genome illuminates the evolution of genome configuration and nuclear architecture of avian chromosomes. *Genome Res*, **31**, 497–511.
- Llopart, A. (2018). Faster-X evolution of gene expression is driven by recessive adaptive cis-regulatory variation in *Drosophila*. *Molecular Ecology*, **27**, 3811–3821.
- Mahadevaraju, S., Fear, J. M., Akeju, M., Galletta, B. J., Pinheiro, M. M. L. S., Avelino, C. C., Cabral-de-Mello, D. C., Conlon, K., Dell’Orso, S., Demere, Z., Mansuria, K., Mendonça, C. A., Palacios-Gimenez, O. M., Ross, E., Savery, M., Yu, K., Smith, H. E., Sartorelli, V., Yang, H., ... Oliver, B. (2021). Dynamic sex chromosome expression in *Drosophila* male germ cells. *Nature Communications*, **12**, 892.
- Mahajan, S., & Bachtrog, D. (2017). Convergent evolution of Y chromosome gene content in flies. *Nature Communications*, **8**, 785.
- Meisel, R. P., & Connallon, T. (2013). The faster-X effect: Integrating theory and data. *Trends in Genetics*, **29**, 537–544.
- Mongue, A. J., Hansen, M. E., & Walters, J. R. (2022). Support for faster and more adaptive Z chromosome evolution in two divergent lepidopteran lineages\*. *Evolution*, **76**, 332–345.
- Muller, H. J. (1914). A gene for the fourth chromosome of *Drosophila*. *Journal of Experimental Zoology*, **17**, 325–336.
- Nei, M. (1970). Accumulation of nonfunctional genes on sheltered chromosomes. *The American Naturalist*, **104**, 311–322.
- Nozawa, M., Minakuchi, Y., Satomura, K., Kondo, S., Toyoda, A., & Tamura, K. (2021). Shared evolutionary trajectories of three independent neo-sex chromosomes in *Drosophila*. *Genome Research*, **31**, 2069–2079.
- Nozawa, M., Onizuka, K., Fujimi, M., Ikeo, K., & Gojobori, T. (2016). Accelerated pseudogenization on the neo-X chromosome in *Drosophila miranda*. *Nature Communications*, **7**, 13659.
- Olito, C., Ponnikas, S., Hansson, B., & Abbott, J. K. (2022). Consequences of partially recessive deleterious genetic variation for the evolution of inversions suppressing recombination between sex chromosomes. *Evolution; International Journal of Organic Evolution*, **76**, 1320–1330.
- Orr, H. A., & Betancourt, A. J. (2001). Haldane’s sieve and adaptation from the standing genetic variation. *Genetics*, **157**, 875–884.
- Patten, M. M. (2019). The X chromosome favors males under sexually antagonistic selection. *Evolution*, **73**, 84–91.
- Pinharanda, A., Rousselle, M., Martin, S. H., Hanly, J. J., Davey, J. W., Kumar, S., Galtier, N. & Jiggins, C. D. (2019). Sexually dimorphic gene expression and transcriptome evolution provide mixed evidence for a fast-Z effect in *Heliconius*. *Journal of Evolutionary Biology*, **32**, 194–204.
- Ponnikas, S., Sigeman, H., Abbott, J. K., & Hansson, B. (2018). Why do sex chromosomes stop recombining? *Trends in Genetics*, **34**, 492–503.
- Presgraves, D. C., & Orr, H. A. (1998). Haldane’s rule in taxa lacking a hemizygous X. *Science*, **282**, 952–954.
- Pucholt, P., Wright, A. E., Conze, L. L., Mank, J. E., & Berlin, S. (2017). Recent sex chromosome divergence despite ancient dioecy in the willow *Salix viminalis*. *Molecular Biology and Evolution*, **34**, 1991–2001.
- Radhakrishnan, S., & Valenzuela, N. (2017). Chromosomal context affects the molecular evolution of sex-linked genes and their autosomal counterparts in turtles and other vertebrates. *Journal of Heredity*, **108**, 720–730.
- Rice, W. R. (1984). Sex chromosomes and the evolution of sexual dimorphism. *Evolution*, **38**, 735–742.
- Rousselle, M., Faivre, N., Ballenghien, M., Galtier, N., & Nabholz, B. (2016). Hemizygoty enhances purifying selection: Lack of fast-Z evolution in two Satyrine butterflies. *Genome Biology and Evolution*, **8**, 3108–3119.
- Rupp, S. M., Webster, T. H., Olney, K. C., Hutchins, E. D., Kusumi, K., & Wilson Sayres, M. A. (2017). Evolution of dosage compensation in *Anolis carolinensis*, a reptile with XX/XY chromosomal sex determination. *Genome Biology and Evolution*, **9**, 231–240.
- Ruzicka, F., & Connallon, T. (2020). Is the X chromosome a hot spot for sexually antagonistic polymorphisms? Biases in current empirical tests of classical theory. *Proceedings of the Royal Society B: Biological Sciences*, **287**, 20201869.
- Ruzicka, F., & Connallon, T. (2022). An unbiased test reveals no enrichment of sexually antagonistic polymorphisms on the human X chromosome. *Proceedings of the Royal Society B: Biological Sciences*, **289**, 20212314.
- Ruzicka, F., Hill, M. S., Pennell, T. M., Flis, I., Ingleby, F. C., Mott, R., Fowler, K., Morrow, E. H. & Reuter, M. (2019). Genome-wide sexually antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies. *PLOS Biology*, **17**, e3000244.
- Sackton, T. B., Corbett-Detig, R. B., Nagaraju, J., Vaishna, L., Arunkumar, K. P., & Hartl, D. L. (2014). Positive selection drives faster-Z evolution in silkworms. *Evolution*, **68**, 2331–2342.
- Sardell, J. M., Josephson, M. P., Dalziel, A. C., Peichel, C. L., & Kirkpatrick, M. (2021). Heterogeneous histories of recombination suppression on stickleback sex chromosomes. *Molecular Biology and Evolution*, **38**, 4403–4418.
- Veltsos, P., Ridout, K. E., Touns, M. A., González-Martínez, S. C., Muyle, A., Emery, O., Rastas, P., Hudzieczek, V., Hobza, R., Vyskot, B., Marais, G. A. B., Filatov, D. A. & Pannell, J. R. (2019). Early sex-chromosome evolution in the diploid dioecious plant *Mercurialis annua*. *Genetics*, **212**, 815–835.
- Vicoso, B. (2019). Molecular and evolutionary dynamics of animal sex-chromosome turnover. *Nature Ecology & Evolution*, **3**, 1632–1641.
- Vicoso, B., & Charlesworth, B. (2006). Evolution on the X chromosome: Unusual patterns and processes. *Nature Reviews. Genetics*, **7**, 645–653.
- Vicoso, B., & Charlesworth, B. (2009). Effective population size and the faster-X effect: An extended model. *Evolution*, **63**, 2413–2426.
- Wright, A. E., Darolti, I., Bloch, N. I., Oostra, V., Sandkam, B., Buechel, S. D., Kolm, N., Breden, F., Vicoso, B. & Mank, J. E. (2017). Convergent recombination suppression suggests role of sexual selection in guppy sex chromosome formation. *Nature Communications*, **8**, 14251.
- Wright, A. E., Dean, R., Zimmer, F., & Mank, J. E. (2016). How to make a sex chromosome. *Nature Communications*, **7**, 12087.
- Zhang, Y. E., Vıbranovski, M. D., Krinsky, B. H., & Long, M. (2010). Age-dependent chromosomal distribution of male-biased genes in *Drosophila*. *Genome Research*, **20**, 1526–1533.
- Zhou, Q., & Bachtrog, D. (2012). Sex-specific adaptation drives early sex chromosome evolution in *Drosophila*. *Science*, **337**, 341–345.