




New Inertial Projection Methods for Solving Multivalued Variational Inequality Problems Beyond Monotonicity

Chinedu Izuchukwu¹ · Yekini Shehu^{2,3} 

Accepted: 13 January 2021 / Published online: 3 March 2021
© The Author(s) 2021

Abstract

In this paper, we present two new inertial projection-type methods for solving multivalued variational inequality problems in finite-dimensional spaces. We establish the convergence of the sequence generated by these methods when the multivalued mapping associated with the problem is only required to be locally bounded without any monotonicity assumption. Furthermore, the inertial techniques that we employ in this paper are quite different from the ones used in most papers. Moreover, based on the weaker assumptions on the inertial factor in our methods, we derive several special cases of our methods. Finally, we present some experimental results to illustrate the profits that we gain by introducing the inertial extrapolation steps.

Keywords Inertial methods · Multivalued variational inequalities · Projection-type methods · Continuous mapping · Armijo-type linesearch

1 Introduction

Assume that C is a nonempty closed and convex subset of \mathbb{R}^N and $F : C \rightrightarrows \mathbb{R}^N$ a multivalued mapping with nonempty values. The Multivalued Variational Inequality

✉ Yekini Shehu
yekini.shehu@unn.edu.ng

Chinedu Izuchukwu
izuchukwu_c@yahoo.com; izuchukwuc@ukzn.ac.za

¹ School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Durban, South Africa

² Department of Mathematics, Zhejiang Normal University, Jinhua 321004, People's Republic of China

³ Institute of Science and Technology (IST), Am Campus 1, 3400, Klosterneuburg, Austria

Problem (MVIP) associated with F and C consists in finding $x^* \in C$ and $u \in F(x^*)$ such that

$$\langle u, y - x^* \rangle \geq 0, \quad \forall y \in C. \quad (1)$$

MVIP (1) was first introduced and studied by Browder (1965) as an important generalization of the classical Variational Inequality Problem (VIP). The MVIP is also known to be a useful generalization of the class of multivalued complementarity problems (see Dong et al. 2017; Facchinei and Pang 2003; He et al. 2019), as well as constrained convex non-smooth optimization problems (see Dong et al. 2017; He et al. 2019; Rockafellar 1970). Therefore, problem (1) is quite general and provides a unified treatment for the study of a wide class of problems such as price equilibrium problems, oligopolistic market equilibrium problems, Nash equilibrium problems, fixed point problems for multivalued mappings, game theory, among others (see Attouch and Cabot 2019b; Brouwer 1912; Carey and Ge 2012; He et al. 2019; Nadler 1969; Oggioni et al. 2012; Raciti and Falsaperla 2007 and the references therein).

When F is a singlevalued mapping in Eq. 1 (i.e., the case of the classical VIP), many methods have been designed by numerous authors for solving the VIP. These include, the gradient projection methods, extragradient methods (Korpelevich 1976), the subgradient extragradient methods (Censor et al. 2011), Tseng's method (Tseng 2000), among others (see, for example, Vuong 2019). We note that these methods for solving the classical VIP are not simply transformed to the case of MVIP since it is very difficult to handle the multivalued mapping associated with the MVIP. Thus, the methods for solving the MVIP are quite different. In 2014, Fang and Chen (2014) extended the subgradient extragradient method for solving the MVIP (1) in finite dimensional spaces. By employing the **Procedure A** below, they proposed the following Algorithm 1.1.

Procedure A (Konnov 1998)

- Input: a point $x \in \mathbb{R}^N$.
 Output: a point $R(x) \in C$, where $C := \{x \in \mathbb{R}^N \mid g(x) \leq 0\}$, $g : \mathbb{R}^N \rightarrow \mathbb{R}$ is a convex function.
 Step 1: set $n = 0$ and $x_n = x$.
 Step 2: if $g(x_n) \leq 0$, then stop and set $R(x) = x_n$. Otherwise, go to Step 3.
 Step 3: choose a point $w_n \in \partial g(x_n)$, where $\partial g(x)$ denotes the subdifferential of g at x , set

$$x_{n+1} = x_n - 2g(x_n) \frac{w_n}{\|w_n\|^2},$$

and set $n := n + 1$ go back to Step 2.

Algorithm 1.1.

- Step 0:** Choose $\bar{x}_1 \in \mathbb{R}^N$ and two parameters $\gamma, \delta \in (0, 1)$. Set $n = 1$.
Step 1. Apply **Procedure A** with $x = \bar{x}_n$ and set $x_n = R(\bar{x}_n)$.

Step 2. Choose $u_n \in F(x_n)$ and let k_n be the smallest nonnegative integer satisfying $v_n \in F(P_C(x_n - \gamma^{k_n} u_n))$,

$$\gamma^{k_n} \|u_n - v_n\| \leq (1 - \delta) \|x_n - P_C(x_n - \gamma^{k_n} u_n)\|. \tag{2}$$

Set $\rho_n = \gamma^{k_n}$ and $z_n = P_C(x_n - \rho_n u_n)$. If $x_n = z_n$, stop.

Step 3. Compute $\bar{x}_{n+1} = P_{C_n}(x_n - \rho_n v_n)$, where $C_n = \{y \in \mathbb{R}^N : \langle x_n - \rho_n u_n - z_n, y - z_n \rangle \leq 0\}$.

Let $n = n + 1$ and return to Step 1.

Inspired by Algorithm 1.1, Dong et al. (2017) proposed the following projection and contraction method for solving the MVIP (1).

Algorithm 1.2.

Step 0: Choose $\bar{x}_1 \in \mathbb{R}^N$ and four parameters $\tau > 0, \gamma, \delta \in (0, 1)$ and $\alpha \in (0, 2)$. Set $n = 1$.

Step 1. Apply **Procedure A** with $x = \bar{x}_n$ and set $x_n = R(\bar{x}_n)$.

Step 2. Choose $u_n \in F(x_n)$ and find the smallest nonnegative integer l_k such that $\rho_n = \tau \gamma^{l_k}$ and $v_n \in F(P_C(x_n - \rho_n u_n))$, which satisfies

$$\rho_n \|u_n - v_n\| \leq (1 - \delta) \|x_n - P_C(x_n - \rho_n u_n)\|. \tag{3}$$

Set $y_n = P_C(x_n - \rho_n u_n)$. If $x_n = y_n$, stop.

Step 3. Compute $\bar{x}_{n+1} = x_n - \alpha \beta_n d(x_n, y_n)$, where $d(x_n, y_n) = \rho_n(u_n - v_n)$, $\phi(x_n, y_n) := \langle x_n - y_n, d(x_n, y_n) \rangle$ and $\beta_n := \frac{\phi(x_n, y_n)}{\|d(x_n, y_n)\|^2}$.

Let $n = n + 1$ and return to Step 1.

We comment that the Armijo-type linesearch procedures (2) and (3) of Algorithm 1.1 and Algorithm 1.2, respectively, involve the computation of projection onto C multiple times in each linesearch. They also involve the evaluation of the multivalued mapping F too many times in each search. To overcome some of these shortcomings, He et al. (2019), proposed the following projection-type method for solving MVIP (1):

Algorithm 1.3.

Step 0: Choose $x_1 \in \mathbb{R}^N$ as an initial point and fix four parameters $\gamma, \sigma \in (0, 1)$ and $0 < \rho^0 \leq \rho^1 < \infty$. Set $C_1 = \mathbb{R}^N, \bar{x}_1 = x_1$, and $n = 1$.

Step 1: Apply **Procedure A** to obtain $x_n = R(\bar{x}_n)$.

Step 2: Choose $u_n \in F(x_n)$ and $\rho_n \in [\rho^0, \rho^1]$. Set $y_n = P_C(x_n - \rho_n u_n)$. If $x_n = y_n$, then stop. Otherwise, compute $z_n = \alpha_n y_n + (1 - \alpha_n) x_n$ and choose the largest $\alpha \in \{\gamma^0, \gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists $w_n \in F(z_n)$ satisfying

$$\langle w_n, x_n - y_n \rangle \geq \sigma \langle u_n, x_n - y_n \rangle.$$

Step 3: Taking a point $v_n \in F(y_n)$, set $d(x_n, y_n) = (x_n - y_n) - \rho_n(u_n - v_n)$ and compute $\bar{v}_n = x_n - \beta_n d(x_n, y_n)$, where $\beta_n = \frac{\phi(x_n, y_n)}{\|d(x_n, y_n)\|^2}$, $\phi(x_n, y_n) = \langle x_n - y_n, d(x_n, y_n) \rangle$.

Step 4: Set $C_n = \{y \in \mathbb{R}^N \mid \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \bigcap_{i=1}^n C_i$.
 Compute $\bar{x}_{n+1} = P_{C_n^*}(\bar{v}_n)$.

If $\bar{x}_{n+1} = x_n$, then stop. Otherwise, let $n := n + 1$ and return Step 1.

As observed in He et al. (2019, Section 4), Algorithm 1.1, Algorithm 1.2 and Algorithm 1.3 do not work well in some settings because of the presence of **Procedure A** in the iterative steps. Hence, the authors in He et al. (2019) proposed the following projection-type method without **Procedure A** for solving MVIP (1), which can be implemented in such settings.

Algorithm 1.4.

Step 0: Choose $x_1 \in \mathbb{R}^N$ as an initial point and fix four parameters $\gamma, \sigma \in (0, 1)$ and $0 < \rho^0 \leq \rho^1 < \infty$. Set $C_1 = \mathbb{R}^N$ and $n = 1$.

Step 1: Choose $u_n \in F(x_n)$ and $\rho_n \in [\rho^0, \rho^1]$. Set $y_n = P_C(x_n - \rho_n u_n)$. If $x_n = y_n$, then stop. Otherwise, compute $z_n = \alpha_n y_n + (1 - \alpha_n)x_n$ and choose the largest $\alpha \in \{\gamma^0, \gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists $w_n \in F(z_n)$ satisfying

$$\langle w_n, x_n - y_n \rangle \geq \sigma \langle u_n, x_n - y_n \rangle.$$

Step 2: Taking a point $v_n \in F(y_n)$, set $d(x_n, y_n) = (x_n - y_n) - \rho_n(u_n - v_n)$ and compute $\bar{x}_n = x_n - \beta_n d(x_n, y_n)$, where $\beta_n = \frac{\phi(x_n, y_n)}{\|d(x_n, y_n)\|^2}$, $\phi(x_n, y_n) = \langle x_n - y_n, d(x_n, y_n) \rangle$.

Step 3: Set $C_n = \{y \in \mathbb{R}^N \mid \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \bigcap_{i=1}^n C_i$.
 Compute $x_{n+1} = P_{C \cap C_n^*}(\bar{x}_n)$.

If $x_{n+1} = x_n$, then stop. Otherwise, let $n := n + 1$ and return Step 1.

Notice that the linesearch procedure in Algorithm 1.3 and Algorithm 1.4 involve the computation of the projection onto C only one time in each search trial. Thus, Algorithm 1.3 and Algorithm 1.4 seem more efficient than Algorithm 1.1 and Algorithm 1.2. Moreover, He et al. (2019) showed numerically that their methods perform better than Algorithm 1.2 of Dong et al. (2017). However, Algorithm 1.3 and Algorithm 1.4 still involve the evaluation of the multivalued mapping at least 3 times in each iteration.

Recently, inertial type algorithms for solving optimization problems have become of great interest to numerous researchers. Since Polyak (1964) studied an inertial extrapolation process for solving the smooth convex minimization problems, there have been growing interests in the design and study of iterative methods with inertial term. For example, inertial forward-backward splitting methods (Attouch et al. 2000; Cholamjiak et al. 2018; Ochs et al. 2015), inertial Douglas-Rachford splitting method (Bot et al. 2015), inertial ADMM (Bot and Csetnek 2016), and inertial forward-backward-forward method (Lorenz and Pock 2015). The inertial term is based upon a discrete analogue of a second order dissipative dynamical system (Attouch et al. 2000) and known for its efficiency in improving the convergence rate of iterative methods. The inertial type algorithms have been tested in the solution of certian number of problems (for example, imaging and data analysis problems, motion of a body

in a potential field) and the tests show that they actually give remarkable speed-up when compared with corresponding algorithms without inertial term (see, for example, Attouch and Cabot 2019a; Attouch and Cabot 2019b; Attouch et al. 2000; Beck and Teboulle 2009; Bot and Csetnek 2016; Lorenz and Pock 2015; Ochs et al. 2015; Polyak 1964; Shehu and Cholakjiak 2019; Shehu et al. 2019; Shehu et al. 2019 and the references therein).

Inspired by this recent trend on inertial extrapolation type methods for solving optimization problems, our aim in this paper is to design some modifications of Algorithms 1.3 and 1.4, together with new inertial extrapolation techniques to solve problem (1). We present two inertial projection-type methods for solving MVIP (1) when the multivalued mapping F is only assumed to be locally bounded without any monotonicity assumption. The first method uses a linesearch as in Algorithm 1.3 and Algorithm 1.4 while the second method uses a different linesearch procedure with the aim of minimizing the number of evaluation of the multivalued mapping F in each search. Furthermore, the inertial techniques that we employ in this paper are quite different from the ones used in most papers (see for example Cholakjiak et al. 2018; Chuang 2017; Lorenz and Pock 2015; Mainge 2008; Moudafi and Oliny 2003; Ochs et al. 2015; Polyak 1964; Shehu and Cholakjiak 2019; Shehu et al. 2019; Shehu et al. 2019; Thong and Hieu 2018; Thong and Hieu 2017 and the references therein). Moreover, based on the weaker assumptions on the inertial factor in our methods, we derive several special cases of our methods. Finally, we provide some numerical implementations of our methods and compare them with the methods in He et al. (2019), in order to show the profits that we gain by introducing the inertial extrapolation steps.

We organize the rest of the paper as follows: We first recall some basic results in Section 2. Some discussions about our methods are given in Section 3. In Section 4, we investigate the convergence analysis of our first method. In Section 5, we analyze the convergence of our second method. In Section 6, we give some numerical experiments to support our theoretical findings. Then, we conclude with some final remarks in Section 7.

2 Preliminaries

The metric projection, denoted by P_C , is a map defined on \mathbb{R}^N onto C which assigns to each $x \in \mathbb{R}^N$, the unique point in C , denoted by P_Cx such that

$$\|x - P_Cx\| = \inf\{\|x - y\| : y \in C\}.$$

It is well known that P_C is nonexpansive, and characterized by the inequality

$$\langle x - P_Cx, y - P_Cx \rangle \leq 0 \quad \forall y \in C. \tag{4}$$

Furthermore, the P_C is known to possess the following property

$$\|P_Cx - x\|^2 \leq \|x - y\|^2 - \|P_Cx - y\|^2 \quad \forall y \in C. \tag{5}$$

It is also known that P_C satisfies

$$\langle x - z, x - P_Cz \rangle \geq \|x - P_Cz\|^2, \quad \forall x \in C, z \in \mathbb{R}^N. \tag{6}$$

For more information and properties of P_C , see Goebel and Reich (1984) and He (2006).

Definition 2.1 A multivalued mapping $F : C \rightrightarrows \mathbb{R}^N$ is said to be

- outer-semicontinuous at $x \in C$ if and only if the graph of F is closed;
- inner-semicontinuous at $x \in C$ if for any sequence $\{x_n\}$ converging to x and $y \in F(x)$, then there exists a sequence $\{y_n\}$ in $F(x_n)$ such that $\{y_n\}$ converges to y ;
- continuous at $x \in C$ if it is both outer-semicontinuous and inner-semicontinuous at x ;
- locally bounded on C if for every $x \in C$, there exists a neighborhood U of x such that $F(U)$ is bounded, where $F(U) = \cup_{x \in U} F(x)$.

Definition 2.2 A multivalued mapping $F : C \rightrightarrows \mathbb{R}^N$ is said to be

- monotone on C if for any $x, y \in C$,

$$\langle u - v, x - y \rangle \geq 0, \quad \forall u \in F(x), v \in F(y);$$

- pseudomonotone on C if for any $x, y \in C$,

$$\text{there exists } u \in F(x) : \langle u, y - x \rangle \geq 0 \text{ implies } \forall v \in F(y) : \langle v, y - x \rangle \geq 0;$$

- quasimonotone on C if for any $x, y \in C$,

$$\text{there exists } u \in F(x) : \langle u, y - x \rangle > 0 \text{ implies } \forall v \in F(y) : \langle v, y - x \rangle \geq 0.$$

Proposition 2.3 (Rockafellar and Wets 2004) *A multivalued mapping $F : C \rightrightarrows \mathbb{R}^N$ is said to be locally bounded if and only if for any bounded sequence $\{x_n\}$ with $u_n \in F(x_n)$, the sequence $\{u_n\}$ is bounded.*

Proposition 2.4 (He et al. 2019) *Assume that the solution set of problem (1) Γ is nonempty and that $F : C \rightrightarrows \mathbb{R}^N$ is continuous. If either*

- F is monotone or pseudomonotone on C ;*
- F is quasimonotone on C and for any $x^* \in \Gamma$ with $u^* \in F(x^*)$ satisfying (1) such that*

$$\text{there exists } y^* \in C : \langle u^*, y^* - x^* \rangle \neq 0;$$

- F is quasimonotone on C with $\text{int } C \neq \emptyset$ and $0 \notin F(x^*)$ for all $x^* \in \Gamma$.*

Then,

$$\langle u, y - x^* \rangle \geq 0 \quad \forall y \in C, u \in F(y), x^* \in \Gamma. \quad (7)$$

Remark 2.5 We can see from Proposition 2.4 that condition (7) is a weaker condition than various monotonicity conditions. Thus, we shall assume for the rest of this paper, that the solution set of problem (1) Γ is nonempty and that Eq. 7 is satisfied.

Following Attouch and Cabot (2019a, pages 5, 10), we note that if $x_{n+1} = x_n + \theta_n(x_n - x_{n-1})$, then for all $n \geq 1$, we have that

$$x_{n+1} - x_n = \left(\prod_{j=1}^n \theta_j \right) (x_1 - x_0),$$

which implies that

$$x_n = x_1 + \left(\sum_{j=1}^{n-1} \prod_{j=1}^l \theta_j \right) (x_1 - x_0).$$

Thus, $\{x_n\}$ converges if and only if $x_1 = x_0$ or if $\sum_{l=1}^{\infty} \prod_{j=1}^l \theta_j < \infty$.

Therefore, we assume henceforth that

$$\sum_{l=i}^{\infty} \left(\prod_{j=i}^l \theta_j \right) < \infty \quad \forall i \geq 1. \tag{8}$$

Then, we can define the sequence $\{t_i\}$ in \mathbb{R} by

$$t_i := \sum_{l=i-1}^{\infty} \left(\prod_{j=i}^l \theta_j \right) = 1 + \sum_{l=i}^{\infty} \left(\prod_{j=i}^l \theta_j \right), \tag{9}$$

with the convention $\prod_{j=i}^{i-1} \theta_j = 1 \quad \forall i \geq 1$.

Remark 2.6 Assumption (8) ensures that $\{t_i\}$ is well-defined in Eq. 9 and

$$t_i = 1 + \theta_i t_{i+1}, \quad \forall i \geq 1. \tag{10}$$

The following proposition provides a criterion for ensuring assumption (8). In fact, this condition makes it possible to cover the usual situations.

Proposition 2.7 (Attouch and Cabot 2019a, Proposition 3.1) *Let $\{\theta_n\}$ be a sequence such that $\theta_n \in [0, 1)$ for every $n \geq 1$. Assume that*

$$\lim_{n \rightarrow \infty} \left(\frac{1}{1 - \theta_{n+1}} - \frac{1}{1 - \theta_n} \right) = c,$$

for some $c \in [0, 1)$. Then, we have

- (i) Condition Eq. 8 holds, and $t_{n+1} \sim \frac{1}{(1-c)(1-\theta_n)}$ as $n \rightarrow \infty$.
- (ii) The equivalence $1 - \theta_n \sim 1 - \theta_{n+1}$ holds true as $n \rightarrow \infty$. Hence, $t_{n+1} \sim t_{n+2}$ as $n \rightarrow \infty$.

Remark 2.8 Example of a sequence satisfying the assumptions of Proposition 2.7 (therefore, satisfying assumption (8)) is $\theta_n = 1 - \frac{\bar{\theta}}{n}$, $\bar{\theta} > 1$.

Clearly,

$$\left(\frac{1}{1 - \theta_{n+1}} - \frac{1}{1 - \theta_n} \right) = \frac{1}{\bar{\theta}}(n + 1) - \frac{1}{\bar{\theta}}n = \frac{1}{\bar{\theta}}.$$

Hence,

$$\lim_{n \rightarrow \infty} \left(\frac{1}{1 - \theta_{n+1}} - \frac{1}{1 - \theta_n} \right) = \frac{1}{\bar{\theta}}.$$

Recall that the above example falls within the setting of Nesterov’s extrapolation methods (for instance, see Attouch and Cabot 2019a; Beck and Teboulle 2009; Chambolle and Dossal 2015, Nesterov 1983).

The corresponding finite sum expression of $\{t_i\}$ is defined for $i, n \geq 1$, by

$$t_{i,n} := \begin{cases} \sum_{l=i-1}^{n-1} \left(\prod_{j=i}^l \theta_j \right) = 1 + \sum_{l=i}^{n-1} \left(\prod_{j=i}^l \theta_j \right), & i \leq n, \\ 0, & \text{otherwise.} \end{cases} \tag{11}$$

In the same manner, we have that $\{t_{i,n}\}$ is well-defined and (see also Attouch and Cabot 2019a)

$$t_{i,n} = 1 + \theta_i t_{i+1,n} \quad \forall i \geq 1, n \geq i + 1. \tag{12}$$

The sequences $\{t_i\}$ and $\{t_{i,n}\}$ are very crucial to our convergence analysis. In fact, their effect can be seen in the following lemma which also plays a crucial role in establishing our convergence results.

Lemma 2.9 (Attouch and Cabot 2019a, page 42, Lemma B.1). *Let $\{a_n\}, \{\theta_n\}$ and $\{w_n\}$ be sequences of real numbers satisfying*

$$a_{n+1} \leq \theta_n a_n + w_n \quad \text{for every } n \geq 1.$$

Assume that $\theta_n \geq 0$ for every $n \geq 1$.

(a) For every $n \geq 1$, we have

$$\sum_{i=1}^n a_i \leq t_{1,n} a_1 + \sum_{i=1}^{n-1} t_{i+1,n} w_i,$$

where the double sequence $\{t_{i,n}\}$ is defined by Eq. 11.

(b) Under Eq. 8, assume that the sequence $\{t_i\}$ defined by Eq. 9 satisfies

$$\sum_{i=1}^{\infty} t_{i+1} [w_i]_+ < \infty. \text{ Then, the series } \sum_{i \geq 1} [a_i]_+ \text{ is convergent, and}$$

$$\sum_{i=1}^{\infty} [a_i]_+ \leq t_1 [a_1]_+ + \sum_{i=1}^{\infty} t_{i+1} [w_i]_+,$$

where $[t]_+ := \max\{t, 0\}$ for any $t \in \mathbb{R}$.

The following lemmas will also be needed in our convergence analysis.

Lemma 2.10 (Facchinei and Pang 2003) *A point $x^* \in \Gamma$ if and only if $x^* = P_C(x^* - \rho u)$ for some $u \in F(x^*)$ and $\rho > 0$.*

Lemma 2.11 (Attouch and Cabot 2019a, page 7, Lemma 2.1). *Let $\{x_n\}$ be a sequence in \mathbb{R}^N , and let $\{\theta_n\}$ be a sequence of real numbers. Given $z \in \mathbb{R}^N$, define the sequence $\{\Gamma_n\}$ by $\Gamma_n := \frac{1}{2}\|x_n - z\|^2$. Then*

$$\Gamma_{n+1} - \Gamma_n - \theta_n(\Gamma_n - \Gamma_{n-1}) = \frac{1}{2}(\theta_n + \theta_n^2)\|x_n - x_{n-1}\|^2 + \langle x_{n+1} - y_n, x_{n+1} - z \rangle - \frac{1}{2}\|x_{n+1} - y_n\|^2, \tag{13}$$

where $y_n = x_n + \theta_n(x_n - x_{n-1})$.

Lemma 2.12 *The following is well-known:*

$$2\langle x, y \rangle = \|x\|^2 + \|y\|^2 - \|x - y\|^2 = \|x + y\|^2 - \|x\|^2 - \|y\|^2 \quad \forall x, y \in \mathbb{R}^N.$$

Lemma 2.13 (Konnov 1998) *The number of iterations in Procedure A is finite and for any given $x \in \mathbb{R}^N$, it holds that*

$$\|R(x) - y\| \leq \|x - y\|, \quad \forall y \in C.$$

3 Proposed Methods

In this section, we present our methods and discuss their features. We begin with the following assumptions under which we obtain our convergence results.

Assumption 3.1 Suppose that the following hold:

- (a) The feasible set C is nonempty, closed and convex subset of \mathbb{R}^N .
- (b) $F : \mathbb{R}^N \rightrightarrows \mathbb{R}^N$ is locally bounded and continuous.
- (c) Γ is nonempty and satisfies condition (7).
- (d) $\theta_n \in [0, 1[$ for all $n \geq 1$ and there exists $\varepsilon \in (0, 1)$ such that for n large enough, we have

$$(1 - \varepsilon)(1 - \theta_{n-1}) \geq \theta_n t_{n+1} \left(1 + \theta_n + [\theta_{n-1} - \theta_n]_+ \right). \tag{14}$$

We now present some criteria that guarantee assumptions (8) and (14).

Proposition 3.1 *Assume that $\{\theta_n\}$ is a nondecreasing sequence that satisfies $\theta_n \in [0, 1[\quad \forall n \geq 1$ with $\lim_{n \rightarrow \infty} \theta_n = \theta$ such that the following condition holds:*

$$1 - 3\theta > 0. \tag{15}$$

Then assumptions (8) and (14) hold.

Proof Observe that $\theta_n \leq \theta \ \forall n \geq 1$. Thus, we have that assumption (8) is satisfied and $t_n = \frac{1}{1-\theta} \ \forall n \geq 1$ (see Attouch and Cabot 2019a). Now, observe that $1 - 3\theta > 0$ implies that $(1 - \theta) > \frac{\theta(1+\theta)}{1-\theta}$. This further implies that there exists $\epsilon \in (0, 1)$ such that

$$(1 - \epsilon)(1 - \theta) \geq \frac{\theta(1 + \theta)}{1 - \theta}. \tag{16}$$

Since $\theta_n \leq \theta \ \forall n \geq 1$, we obtain from Eq. 16 that

$$(1 - \epsilon)(1 - \theta_{n-1}) \geq \frac{\theta(1 + \theta)}{1 - \theta} \geq \theta_n t_{n+1} (1 + \theta_n), \tag{17}$$

for some $\epsilon \in (0, 1)$. Since $\theta_{n-1} \leq \theta_n \ \forall n \geq 1$, we obtain that

$$\theta_n t_{n+1} (1 + \theta_n) = \theta_n t_{n+1} (1 + \theta_n + [\theta_{n-1} - \theta_n]_+).$$

Combining this with Eq. 17, we get that the assumption (14) is satisfied. □

Proposition 3.2 *Suppose that $\theta_n \in [0, 1)$ and there exists $c \in [0, \frac{1}{2})$ such that*

$$\lim_{n \rightarrow \infty} \left(\frac{1}{1 - \theta_{n+1}} - \frac{1}{1 - \theta_n} \right) = c \tag{18}$$

and

$$\liminf_{n \rightarrow \infty} (1 - \theta_n)^2 > \limsup_{n \rightarrow \infty} \frac{\theta_n(1 + \theta_n)}{1 - 2c}. \tag{19}$$

Then assumption (14) holds.

Proof From Eq. 19, we obtain that

$$\liminf_{n \rightarrow \infty} (1 - \theta_{n-1})^2 > \limsup_{n \rightarrow \infty} \frac{\theta_n(1 + \theta_n)}{1 - 2c}. \tag{20}$$

Thus, there exists $\epsilon \in (0, 1)$ sufficiently small enough such that

$$\liminf_{n \rightarrow \infty} (1 - \theta_{n-1})^2 > \limsup_{n \rightarrow \infty} \frac{\theta_n(1 + \theta_n)}{1 - 2c - \epsilon(1 - c)} > \limsup_{n \rightarrow \infty} \frac{\theta_n(1 + \theta_n)}{1 - 2c}. \tag{21}$$

This implies that

$$\begin{aligned} (1 + o(1))\theta_n(1 + \theta_n) &\leq [1 - 2c - \epsilon(1 - c) + o(1)](1 - \theta_{n-1})^2 \\ &= [(1 - \epsilon)(1 - c) - (2c - c + o(1))](1 - \theta_{n-1})^2 \\ &\leq [(1 - \epsilon)(1 - c) - \theta_n(c + o(1))](1 - \theta_{n-1})^2, \end{aligned}$$

which implies that

$$(1 - \epsilon)(1 - c)(1 - \theta_{n-1})^2 \geq (1 + o(1))\theta_n (1 + \theta_n + (1 - \theta_{n-1})^2 + o((1 - \theta_{n-1})^2)). \tag{22}$$

Now, observe from Eq. 18 that

$$\theta_{n-1} - \theta_n + c(1 - \theta_{n-1})(1 - \theta_n) = o((1 - \theta_{n-1})(1 - \theta_n)),$$

which implies from Proposition 2.7(ii) that

$$\begin{aligned} \theta_{n-1} - \theta_n &= -c(1 - \theta_{n-1})(1 - \theta_n) + o((1 - \theta_{n-1})(1 - \theta_n)) \\ &= -c(1 - \theta_{n-1})^2 + o(1 - \theta_{n-1})^2 \text{ as } n \rightarrow \infty. \end{aligned}$$

This implies that

$$\begin{aligned} |\theta_{n-1} - \theta_n| &= | -c(1 - \theta_{n-1})^2 + o(1 - \theta_{n-1})^2 | \\ &\leq c(1 - \theta_{n-1})^2 + o(1 - \theta_{n-1})^2 \text{ as } n \rightarrow \infty. \end{aligned} \tag{23}$$

Combining (22) and (23), we obtain that

$$(1 - \epsilon)(1 - c)(1 - \theta_{n-1})^2 \geq (1 + o(1))\theta_n (1 + \theta_n + [\theta_{n-1} - \theta_n]_+). \tag{24}$$

By Proposition 2.7, we have that $t_{n+1} \sim t_n \sim \frac{1}{(1-c)(1-\theta_{n-1})}$ as $n \rightarrow \infty$. Hence, Eq. 24 is equivalent to

$$(1 - \epsilon)(1 - c)(1 - \theta_{n-1})^2 \geq \frac{\theta_n}{(1-c)(1-\theta_{n-1})} t_{n+1} (1 + \theta_n + [\theta_{n-1} - \theta_n]_+),$$

which implies that assumption (14) holds. □

Remark 3.3 We mention that Proposition 3.1 and Proposition 3.2 provide some sufficient conditions for ensuring that assumptions (14) and (8) hold. That is, assumptions (14) and (8) are much more weaker conditions than the assumptions in both propositions. Note that similar conditions as in Propositions 3.1 and 3.2 have been used by other authors to ensure convergence of inertial methods (see Lorenz and Pock 2015; Thong and Hieu 2018; Thong and Hieu 2017 and the references therein). In fact, we shall see later that using the conditions in Proposition 3.1 and Proposition 3.2, we derive some corollaries of our results.

We now present the first method of this paper.

Algorithm 3.2.

Step 0: Choose the sequence $\{\theta_n\}$ in $[0, 1)$ such that the condition from Eqs. 8 and 14 hold. Let $x_1, x_0 \in \mathbb{R}^N$ be given arbitrary and fix $\gamma, \sigma \in (0, 1), 0 < \rho_0 \leq \rho_1 < \infty$. Set $C_1 = \mathbb{R}^N$ and $n = 1$.

Step 1. Set

$$v_n = x_n + \theta_n(x_n - x_{n-1})$$

and choose $u_n \in F(v_n)$ and $\rho_n \in [\rho_0, \rho_1]$. Then, compute

$y_n = P_C(v_n - \rho_n u_n)$. If $v_n = y_n$: STOP. Otherwise, go to **Step 2**.

Step 2. Compute

$$z_n = \alpha_n y_n + (1 - \alpha_n)v_n$$

and choose the largest $\alpha \in \{\gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists a point $w_n \in F(z_n)$ satisfying

$$\langle w_n, v_n - y_n \rangle \geq \sigma \langle u_n, v_n - y_n \rangle. \tag{25}$$

Step 3. Set $C_n = \{y \in \mathbb{R}^N : \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \bigcap_{i=1}^n C_i$. Then, compute

$$x_{n+1} = P_{C_n^*}(v_n).$$

Set $n := n + 1$ and go back to **Step 1**.

Lemma 3.4 *Step 2 of Algorithm 3.2 is well-defined.*

Proof Let $v \in C$ and $u \in F(v)$. Define $y := P_C(v - \rho u)$, $\rho > 0$. If $v = y$, then by Lemma 2.10, we have that v is a solution. Now, if $v \neq y$, then by Eq. 4,

$$\langle u, v - y \rangle = \frac{1}{\rho} \langle y - (v - \rho u) + (v - y), v - y \rangle \geq \frac{1}{\rho} \langle v - y, v - y \rangle > 0. \tag{26}$$

Now, suppose on the contrary that Step 2 is not well-defined, then we will have that, for any $\alpha > 0$ and $w \in F(z)$ with $z = \alpha y + (1 - \alpha)v$,

$$\langle w, v - y \rangle < \sigma \langle u, v - y \rangle. \tag{27}$$

In particular, for $\alpha_n = \frac{1}{n^2}$ with $z_n = \alpha_n y + (1 - \alpha_n)v$, we have that $z_n \rightarrow v$ as $n \rightarrow \infty$. Since F is continuous, it is inner-semicontinuous. Thus, there exists $w_n \in F(z_n)$ such that $w_n \rightarrow u$ with $u \in F(v)$. Taking w as w_n in Eq. 27, and taking limit as $n \rightarrow \infty$, we obtain that

$$(1 - \sigma) \langle u, v - y \rangle \leq 0,$$

which contradicts (26). Hence, Step 2 of Algorithm 3.2 is well-defined. □

Remark 3.5 Observe that Assumption 3.1 (c) ensures that Step 3 of Algorithm 3.2 is well-defined since $\Gamma \subset C_n^*$ and hence $C_n^* \neq \emptyset$ for all $n \geq 2$. Indeed, for $z \in \Gamma$, we obtain from Assumption 3.1 (c) that $\langle w_n, z - z_n \rangle \leq 0 \forall n \geq 2$. Thus, $z \in C_n \forall n \geq 2$, which follows that $z \in C_n^* \forall n \geq 2$.

In the following, we present another method with a new linesearch (different from Eq. 25) with the aim of minimizing the number of evaluation of the multivalued mapping F in each search.

Algorithm 3.3.

Step 0: Choose the sequence $\{\theta_n\}$ such that the condition from Eqs. 8 and 14 hold. Let $x_1, x_0 \in \mathbb{R}^N$ be given arbitrary and fix $\gamma, \sigma \in (0, 1), 0 < \rho_0 \leq \rho_1 < \infty$. Set $C_1 = \mathbb{R}^N$ and $n = 1$.

Step 1. Set

$$v_n = x_n + \theta_n(x_n - x_{n-1})$$

and choose $u_n \in F(v_n)$ and $\rho_n \in [\rho_0, \rho_1]$. Then, compute

$$y_n = P_C(v_n - \rho_n u_n). \text{ If } v_n = y_n: \text{ STOP. Otherwise, go to Step 2.}$$

Step 2. Compute

$$z_n = \alpha_n y_n + (1 - \alpha_n)v_n$$

and choose the largest $\alpha \in \{\gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists a point $w_n \in F(z_n)$ satisfying

$$\langle w_n, v_n - y_n \rangle \geq \frac{\sigma}{2} \|v_n - y_n\|^2.$$

Step 3. Set $C_n = \{y \in \mathbb{R}^N : \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \cap_{i=1}^n C_i$. Then, compute

$$x_{n+1} = P_{C_n^*}(v_n).$$

Set $n := n + 1$ and go back to **Step 1**.

Remark 3.6 (a) Observe that if we choose a point $u \in F(x)$ with $y := P_C(x - \rho u)$, then, by setting $z = x - \rho u$ in Eq. 6, we obtain that

$$\langle u, x - y \rangle \geq \frac{\sigma}{2} \|x - y\|^2. \tag{28}$$

Thus, using Eq. 28 and the continuity of F , we can see that Step 2 of Algorithm 3.3 is well-defined.

(b) Our Algorithm 3.2 and Algorithm 3.3 have fewer evaluations of multivalued mapping F than Algorithm 1.3 and Algorithm 1.4.

4 Convergence Analysis for Algorithm 3.2

Lemma 4.1 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2 and $\{\Gamma_n\}$ be defined by $\Gamma_n = \frac{1}{2} \|x_n - z\|^2$ for any $z \in \Gamma$. Then, under assumption (8) and Assumption 3.1(c),(d), we have that*

$$\begin{aligned} & \sum_{i=1}^{n-1} [t_{i+1,n} ((1 - 3\theta_i) - (1 - \theta_i)) + t_{i,n}(1 - \theta_{i-1})] \|x_i - x_{i-1}\|^2 \\ & \leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0 + t_1(1 - \theta_0)\|x_1 - x_0\|^2, \end{aligned}$$

where $\{t_{i,n}\}$ is defined in Eq. 11.

Proof First observe that

$$\begin{aligned} \|x_{n+1} - x_n\|^2 &= \|x_{n+1} - 2x_n + x_{n-1} - (x_{n-1} - x_n)\|^2 \\ &= \|x_{n+1} - 2x_n + x_{n-1}\|^2 + \|x_{n-1} - x_n\|^2 \\ &\quad + 2\langle x_{n+1} - 2x_n + x_{n-1}, x_n - x_{n-1} \rangle, \end{aligned}$$

which implies that

$$2\langle x_{n+1} - 2x_n + x_{n-1}, x_n - x_{n-1} \rangle = \|x_{n+1} - x_n\|^2 - \|x_{n+1} - 2x_n + x_{n-1}\|^2 - \|x_{n-1} - x_n\|^2.$$

Thus, we obtain that

$$\begin{aligned} \|x_{n+1} - v_n\|^2 &= \|x_{n+1} - x_n - (x_n - x_{n-1}) + (1 - \theta_n)(x_n - x_{n-1})\|^2 \\ &= \|x_{n+1} - 2x_n + x_{n-1}\|^2 + (1 - \theta_n)^2 \|x_n - x_{n-1}\|^2 \\ &\quad + 2(1 - \theta_n)\langle x_{n+1} - 2x_n + x_{n-1}, x_n - x_{n-1} \rangle \\ &= \|x_{n+1} - 2x_n + x_{n-1}\|^2 + (1 - \theta_n)^2 \|x_n - x_{n-1}\|^2 \\ &\quad + (1 - \theta_n) \left[\|x_{n+1} - x_n\|^2 - \|x_n - x_{n-1}\|^2 - \|x_{n+1} - 2x_n + x_{n-1}\|^2 \right] \\ &= \theta_n \|x_{n+1} - 2x_n + x_{n-1}\|^2 + (1 - \theta_n)^2 \|x_n - x_{n-1}\|^2 \\ &\quad + (1 - \theta_n) \left[\|x_{n+1} - x_n\|^2 - \|x_n - x_{n-1}\|^2 \right] \\ &\geq (1 - \theta_n)^2 \|x_n - x_{n-1}\|^2 + (1 - \theta_n) \left[\|x_{n+1} - x_n\|^2 - \|x_n - x_{n-1}\|^2 \right]. \end{aligned} \tag{29}$$

Let $z \in \Gamma$, then by Remark 3.5, we have that $z \in C_n^*$. Thus, we obtain from Lemma 2.11 and Eq. 29 that

$$\begin{aligned} \Gamma_{n+1} - \Gamma_n - \theta_n(\Gamma_n - \Gamma_{n-1}) &= \frac{1}{2}(\theta_n + \theta_n^2)\|x_n - x_{n-1}\|^2 + \langle x_{n+1} - v_n, x_{n+1} - z \rangle \\ &\quad - \frac{1}{2}\|x_{n+1} - v_n\|^2 \\ &\leq \frac{1}{2}(\theta_n + \theta_n^2)\|x_n - x_{n-1}\|^2 - \frac{1}{2}\|x_{n+1} - v_n\|^2 \tag{30} \\ &\leq \frac{1}{2}(\theta_n + \theta_n^2)\|x_n - x_{n-1}\|^2 - \frac{1}{2}(1 - \theta_n)^2\|x_n - x_{n-1}\|^2 \\ &\quad - \frac{1}{2}(1 - \theta_n) \left[\|x_{n+1} - x_n\|^2 - \|x_n - x_{n-1}\|^2 \right] \\ &= \frac{1}{2}(3\theta_n - 1)\|x_n - x_{n-1}\|^2 - \frac{1}{2}(1 - \theta_n) \\ &\quad \times \left[\|x_{n+1} - x_n\|^2 - \|x_n - x_{n-1}\|^2 \right], \end{aligned}$$

which implies from Lemma 2.9 (a) that

$$\begin{aligned} \Gamma_n - \Gamma_0 &= \sum_{i=1}^n (\Gamma_i - \Gamma_{i-1}) \\ &\leq t_{1,n} (\Gamma_1 - \Gamma_0) + \sum_{i=1}^{n-1} t_{i+1,n} \left[\frac{1}{2}(3\theta_i - 1)\|x_i - x_{i-1}\|^2 - \frac{1}{2}(1 - \theta_i) \right. \\ &\quad \left. \times \left(\|x_{i+1} - x_i\|^2 - \|x_i - x_{i-1}\|^2 \right) \right]. \end{aligned}$$

Notice that $t_{1,n} \leq t_1$. Thus, we obtain that

$$\begin{aligned} & \sum_{i=1}^{n-1} t_{i+1,n} \left[(1 - 3\theta_i) \|x_i - x_{i-1}\|^2 + (1 - \theta_i) \left(\|x_{i+1} - x_i\|^2 - \|x_i - x_{i-1}\|^2 \right) \right] \\ & \leq 2t_{1,n}(\Gamma_1 - \Gamma_0) + 2(\Gamma_0 - \Gamma_n) \\ & \leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0. \end{aligned} \tag{31}$$

□

Now, observe that

$$\begin{aligned} & \sum_{i=1}^{n-1} t_{i+1,n}(1 - \theta_i) \left(\|x_{i+1} - x_i\|^2 - \|x_i - x_{i-1}\|^2 \right) \\ & = \sum_{i=1}^{n-1} (t_{i,n}(1 - \theta_{i-1}) - t_{i+1,n}(1 - \theta_i)) \|x_i - x_{i-1}\|^2 \\ & \quad + t_{n,n}(1 - \theta_{n-1}) \|x_n - x_{n-1}\|^2 - t_{1,n}(1 - \theta_0) \|x_1 - x_0\|^2 \\ & \geq \sum_{i=1}^{n-1} (t_{i,n}(1 - \theta_{i-1}) - t_{i+1,n}(1 - \theta_i)) \|x_i - x_{i-1}\|^2 - t_1(1 - \theta_0) \|x_1 - x_0\|^2. \end{aligned} \tag{32}$$

Combining (31) and (32), we get that

$$\begin{aligned} & \sum_{i=1}^{n-1} t_{i+1,n}(1 - 3\theta_i) \|x_i - x_{i-1}\|^2 + \sum_{i=1}^{n-1} (t_{i,n}(1 - \theta_{i-1}) - t_{i+1,n}(1 - \theta_i)) \|x_i - x_{i-1}\|^2 \\ & \leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0 + t_1(1 - \theta_0) \|x_0 - x_1\|^2. \end{aligned}$$

That is,

$$\begin{aligned} & \sum_{i=1}^{n-1} [t_{i+1,n} ((1 - 3\theta_i) - (1 - \theta_i)) + t_{i,n}(1 - \theta_{i-1})] \|x_i - x_{i-1}\|^2 \\ & \leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0 + t_1(1 - \theta_0) \|x_0 - x_1\|^2. \end{aligned} \tag{33}$$

Lemma 4.2 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2. Then, under assumption (8) and Assumption 3.1(c),(d), we have that $\sum_{n=1}^{\infty} (1 - \theta_{n-1}) \|x_n - x_{n-1}\|^2 < \infty$*

and $\sum_{n=1}^{\infty} \theta_n t_{n+1} \|x_n - x_{n-1}\|^2 < \infty$.

Proof From Eq. 12 and since $t_{i+1,n} \leq t_{i+1}$, we obtain

$$\begin{aligned}
 & t_{i+1,n} [(1 - 3\theta_i) - (1 - \theta_i)] + t_{i,n}(1 - \theta_{i-1}) \\
 &= t_{i+1,n} [(1 - 3\theta_i) - (1 - \theta_i)] + (1 - \theta_{i-1}) + \theta_i t_{i+1,n}(1 - \theta_{i-1}) \\
 &= t_{i+1,n} [(1 - 3\theta_i) - (1 - \theta_i) + \theta_i(1 - \theta_{i-1})] + (1 - \theta_{i-1}) \\
 &= (1 - \theta_{i-1}) - \theta_i t_{i+1,n} (1 - \theta_{i-1}) \\
 &\geq (1 - \theta_{i-1}) - \theta_i t_{i+1} (1 - \theta_{i-1}) \\
 &\geq (1 - \theta_{i-1}) - \theta_i t_{i+1} \left(1 + \theta_i + [\theta_{i-1} - \theta_i]_+\right). \tag{34}
 \end{aligned}$$

Using Eq. 34 in Lemma 4.1, we obtain that

$$\begin{aligned}
 & \sum_{i=1}^{n-1} (1 - \theta_{i-1}) - \theta_i t_{i+1} \left(1 + \theta_i + [\theta_{i-1} - \theta_i]_+\right) \|x_i - x_{i-1}\|^2 \\
 &\leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0 + t_1(1 - \theta_0)\|x_0 - x_1\|^2.
 \end{aligned}$$

We may assume without loss of generality that assumption (14) holds for every $n \geq 1$. Then, we obtain that

$$\sum_{i=1}^{n-1} \varepsilon(1 - \theta_{i-1})\|x_i - x_{i-1}\|^2 \leq 2t_1|\Gamma_1 - \Gamma_0| + 2\Gamma_0 + t_1(1 - \theta_0)\|x_0 - x_1\|^2.$$

Now, taking limit as $n \rightarrow \infty$, we get that

$$\sum_{i=1}^{\infty} (1 - \theta_{i-1})\|x_i - x_{i-1}\|^2 < \infty. \tag{35}$$

Thus, the first conclusion of the lemma is established. To establish the second conclusion of the lemma, we employ assumption (14) again in Eq. 35 and obtain

$$\sum_{i=1}^{\infty} \theta_i t_{i+1} \|x_i - x_{i-1}\|^2 < \infty.$$

□

Lemma 4.3 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2. Then, under assumption (8) and Assumption 3.1(c),(d), we have that*

- (a) $\lim_{n \rightarrow \infty} \|x_n - z\|$ exists for all $z \in \Gamma$.
- (b) $\lim_{n \rightarrow \infty} \|v_n - x_{n+1}\| = 0$.

Proof (a) From Eq. 30, we obtain that

$$\begin{aligned}
 \Gamma_{n+1} - \Gamma_n &\leq \theta_n(\Gamma_n - \Gamma_{n-1}) + \frac{1}{2}(\theta_n + \theta_n^2)\|x_n - x_{n-1}\|^2 - \frac{1}{2}\|x_{n+1} - v_n\|^2 \\
 &\leq \theta_n(\Gamma_n - \Gamma_{n-1}) + \theta_n\|x_n - x_{n-1}\|^2 - \frac{1}{2}\|x_{n+1} - v_n\|^2 \tag{36} \\
 &\leq \theta_n(\Gamma_n - \Gamma_{n-1}) + \theta_n\|x_n - x_{n-1}\|^2.
 \end{aligned}$$

Thus, from Lemma 4.2 and Lemma 2.9 (b), we obtain that $\sum_{n=1}^{\infty} [\Gamma_n - \Gamma_{n-1}]_+ < \infty$. This implies that $\lim_{n \rightarrow \infty} \Gamma_n = \lim_{n \rightarrow \infty} \frac{1}{2} \|x_n - z\|^2$ exists, which further gives that $\lim_{n \rightarrow \infty} \|x_n - z\|$ exists for all $z \in \Gamma$.

(b) Now, using Eq. 36 and Lemma 2.9 (a), we obtain that

$$\begin{aligned} \Gamma_n - \Gamma_0 &= \sum_{i=1}^n (\Gamma_i - \Gamma_{i-1}) \\ &\leq t_{1,n} (\Gamma_1 - \Gamma_0) + \sum_{i=1}^{n-1} t_{i+1,n} \left[\theta_i \|x_i - x_{i-1}\|^2 - \frac{1}{2} \|x_{i+1} - v_i\|^2 \right]. \end{aligned} \tag{37}$$

Since $t_{i+1,n} \leq t_{i+1}$, we obtain from Eq. 37 and Lemma 4.2 that

$$\begin{aligned} \sum_{i=1}^{n-1} t_{i+1,n} \|x_{i+1} - v_i\|^2 &\leq 2\Gamma_0 + 2t_{1,n} (\Gamma_1 - \Gamma_0) + 2 \sum_{i=1}^{n-1} t_{i+1,n} \theta_i \|x_i - x_{i-1}\|^2 \\ &\leq 2\Gamma_0 + 2t_1 |\Gamma_1 - \Gamma_0| + 2 \sum_{i=1}^{\infty} t_{i+1} \theta_i \|x_i - x_{i-1}\|^2 < \infty. \end{aligned}$$

Since $t_{i+1,n} = 0$ for $i \geq n$, letting n tend to ∞ , we obtain that

$$\sum_{i=1}^{\infty} t_{i+1} \|x_{i+1} - v_i\|^2 < \infty. \tag{38}$$

Replacing i with n in Eq. 38 and since $t_n \geq 1$ for every $n \geq 1$, we obtain from Eq. 38 that $\sum_{n=1}^{\infty} \|x_{n+1} - v_n\|^2 < \infty$. This implies that $\lim_{n \rightarrow \infty} \|v_n - x_{n+1}\| = 0$. □

Remark 4.4 The main role of assumption (14) is to guarantee the condition

$$\sum_{n=1}^{\infty} t_{n+1} \theta_n \|x_n - x_{n-1}\|^2 < \infty, \tag{39}$$

obtained in Lemma 4.2 above. Note that Lemma 4.3 holds true if we assume condition (39) directly. Moreover, if $\theta_n \in [0, \theta]$ for every $n \geq 1$, where $\theta \in [0, 1)$, then $t_n \leq \frac{1}{(1-\theta)}$ $\forall n \geq 1$. Under this setting, we have that condition (39) is guaranteed by the condition

$$\sum_{n=1}^{\infty} \theta_n \|x_n - x_{n-1}\|^2 < \infty. \tag{40}$$

In other words, if we assume that condition (40) holds for $\theta_n \in [0, \theta] \forall n \geq 1$, with $\theta \in [0, 1)$, then Lemma 4.3 holds. This assumption has been used by numerous authors to ensure convergence of inertial methods (see, for example, Alvarez and Attouch 2001; Chuang 2017; Lorenz and Pock 2015; Mainge 2008; Moudafi and Oliny 2003 and the references therein).

Furthermore, under the assumptions of Proposition 3.1, we obtain the following as corollaries of Lemma 4.2 and Lemma 4.3 respectively.

Corollary 4.5 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2 such that Assumption 3.1(c) holds. Suppose that $\{\theta_n\}$ is a nondecreasing sequence that satisfies $\theta_n \in [0, 1[\forall n \geq 1$ with $\lim_{n \rightarrow \infty} \theta_n = \theta$ such that $1 - 3\theta > 0$. Then, we have that*

$$\sum_{n=1}^{\infty} (1 - \theta_{n-1}) \|x_n - x_{n-1}\|^2 < \infty \text{ and } \sum_{n=1}^{\infty} \theta_n t_{n+1} \|x_n - x_{n-1}\|^2 < \infty.$$

Proof By Proposition 3.1, we have that assumptions (8) and (14) hold. Hence, the proof follows from Lemma 4.2. \square

Corollary 4.6 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2 such that Assumption 3.1(c) holds. Suppose that $\{\theta_n\}$ is a nondecreasing sequence that satisfies $\theta_n \in [0, 1[\forall n \geq 1$ with $\lim_{n \rightarrow \infty} \theta_n = \theta$ such that $1 - 3\theta > 0$. Then,*

- (a) $\lim_{n \rightarrow \infty} \|x_n - z\|$ exists for all $z \in \Gamma$.
- (b) $\lim_{n \rightarrow \infty} \|v_n - x_{n+1}\| = 0$.

Proof It is similar to the proof of Corollary 4.5. \square

Remark 4.7 Observe that Eq. 18 and Proposition 2.7 imply that condition (8) also holds in Proposition 3.2. Hence, by replacing assumptions (8) and (14) with the assumptions of Proposition 3.2 in Lemma 4.2 and Lemma 4.3, we also obtain corollaries of Lemma 4.2 and Lemma 4.3 in the same manner as Corollaries 4.5 and 4.6 respectively.

Remark 4.8 If we take the inertial factor θ_n to be a constant (that is $\theta_n = \theta \forall n \geq 1$), then we obtain the following corollaries of Lemma 4.2 and Lemma 4.3.

Corollary 4.9 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2 such that Assumption 3.1(c) holds. Suppose that $\theta_n = \theta \forall n \geq 1$ with $\theta \in [0, 1)$ such that*

$$(1 - \theta)^2 > \theta(1 + \theta). \quad (41)$$

Then, we have that $\sum_{n=1}^{\infty} (1 - \theta) \|x_n - x_{n-1}\|^2 < \infty$ and $\sum_{n=1}^{\infty} \frac{\theta}{1 - \theta} \|x_n - x_{n-1}\|^2 < \infty$.

Consequently, we have $\sum_{n=1}^{\infty} \|x_n - x_{n-1}\|^2 < \infty$.

Proof Since $\theta_n = \theta \in [0, 1)$, we obtain for $i \geq 1$ that $t_i = \sum_{l=i-1}^{\infty} \theta^{l-i+1} = \frac{1}{1-\theta} < \infty$. Thus, we get that assumption (8) holds. Note also from Eq. 41 that there exists $\epsilon \in (0, 1)$ such that

$$(1 - \epsilon)(1 - \theta) \geq \frac{\theta(1 + \theta)}{1 - \theta},$$

which is equivalent to condition (14) since $\theta_n = \theta \quad \forall n \geq 1$. Hence, all the assumptions of Lemma 4.2 are satisfied. Thus, the rest of the proof follows from Lemma 4.2. □

Corollary 4.10 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2 such that Assumption 3.1(c) holds. Suppose that $\theta_n = \theta \quad \forall n \geq 1$ with $\theta \in [0, 1)$ such that $(1 - \theta)^2 > \theta(1 + \theta)$. Then,*

- (a) $\lim_{n \rightarrow \infty} \|x_n - z\|$ exists for all $z \in \Gamma$.
- (b) $\lim_{n \rightarrow \infty} \|v_n - x_{n+1}\| = 0$.

Proof The proof is similar to the proof of Corollary 4.9. □

We now return to a very important result for our convergence analysis, whose proof rely on the linesearch given in Algorithm 3.2.

Lemma 4.11 *Let assumption (8) and Assumption 3.1 hold, and let the sequence $\{x_n\}$ be generated by Algorithm 3.2. Then, $\lim_{n \rightarrow \infty} \alpha_n \|y_n - v_n\|^2 = 0$. Moreover, if there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $\{x_{n_k}\}$ converges to x^* and $x^* \notin \Gamma$, then*

- (a) $\liminf_{k \rightarrow \infty} \alpha_{n_k} > 0$;
- (b) $\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0$.

Proof From Eq. 4, Step 1, Step 2 and the fact that $x_{n+1} \in C_n^*$, we obtain that

$$\begin{aligned} \alpha_n \|v_n - y_n\|^2 &= \alpha_n \langle v_n - y_n, v_n - y_n \rangle \\ &\leq \alpha_n \langle v_n - y_n, v_n - y_n \rangle + \alpha_n \langle y_n - v_n + \rho_n u_n, v_n - y_n \rangle \\ &= \alpha_n \rho_n \langle u_n, v_n - y_n \rangle \\ &\leq \frac{\alpha_n \rho_n}{\sigma} \langle w_n, v_n - y_n \rangle \\ &= \frac{\rho_n}{\sigma} \langle w_n, v_n - z_n \rangle \\ &\leq \frac{\rho_n}{\sigma} (\langle w_n, v_n - x_{n+1} \rangle + \langle w_n, x_{n+1} - z_n \rangle) \\ &\leq \frac{\rho_n}{\sigma} \|w_n\| \|v_n - x_{n+1}\|. \end{aligned} \tag{42}$$

Since by Lemma 4.3, $\{x_n\}$ is bounded, we have that $\{z_n\}$ is also bounded. Moreover, since F is locally bounded, we obtain from Proposition 2.3 that $\{w_n\}$ is bounded. Using this and the boundedness of $\{\rho_n\}$, we obtain from Eq. 42 and Lemma 4.3 that

$$\lim_{n \rightarrow \infty} \alpha_n \|y_n - v_n\|^2 = 0. \tag{43}$$

(a) By Step 2, we have that $\{\alpha_n\} \subset (0, 1)$ is bounded. Thus, there exists a subsequence $\{\alpha_{n_k}\}$ of $\{\alpha_n\}$ such that $\liminf_{k \rightarrow \infty} \alpha_{n_k} \geq 0$.

In fact, we claim that $\liminf_{k \rightarrow \infty} \alpha_{n_k} > 0$. Suppose on the contrary that $\liminf_{k \rightarrow \infty} \alpha_{n_k} = 0$. Then, without loss of generality, we can choose a subsequence of $\{\alpha_{n_k}\}$ still denoted by $\{\alpha_{n_k}\}$ such that $\lim_{k \rightarrow \infty} \alpha_{n_k} = 0$.

Now, define $\bar{\alpha}_{n_k} := \frac{\alpha_{n_k}}{\gamma}$, $\bar{z}_{n_k} := \bar{\alpha}_{n_k} y_{n_k} + (1 - \bar{\alpha}_{n_k}) v_{n_k}$. Then, by the boundedness of $\{y_{n_k} - v_{n_k}\}$ and since $\alpha_{n_k} \rightarrow 0$ as $k \rightarrow \infty$, we obtain that

$$\lim_{k \rightarrow \infty} \|\bar{z}_{n_k} - v_{n_k}\| = 0. \tag{44}$$

Also, by Lemma 4.2, we obtain that $\lim_{k \rightarrow \infty} \|x_{n_k} - v_{n_k}\| = \lim_{k \rightarrow \infty} \theta_{n_k} \|x_{n_k} - x_{n_k-1}\| = 0$. Thus, since $x_{n_k} \rightarrow x^*$, we have that $v_{n_k} \rightarrow x^*$. Using Assumption 3.1 (b), the boundedness of $\{v_{n_k}\}$ and Proposition 2.3, we obtain that $\{u_{n_k}\}$ is also bounded. Thus, we can choose a subsequence of $\{u_{n_k}\}$ still denoted by $\{u_{n_k}\}$ such that $u_{n_k} \rightarrow \bar{u}$. Since F is continuous, it is outer-semicontinuous. Hence, $\bar{u} \in F(x^*)$. We also assume without loss of generality that $\rho_{n_k} \rightarrow \rho \in [\rho_0, \rho_1]$. Therefore, we obtain from the continuity of P_C that $y_{n_k} \rightarrow y^*$ as $k \rightarrow \infty$, where $y^* = P_C(x^* - \rho \bar{u})$.

Again, from Eq. 44, we obtain that $\bar{z}_{n_k} \rightarrow x^*$. Since F is inner-semicontinuous and $\bar{u} \in F(x^*)$, we can choose a subsequence $w_{n_k} \in F(\bar{z}_{n_k})$ such that $\bar{w}_{n_k} \rightarrow \bar{u}$.

Now, from the definition of \bar{z}_{n_k} and Step 2, we obtain that

$$\langle \bar{w}_{n_k}, v_{n_k} - y_{n_k} \rangle < \sigma \langle u_{n_k}, v_{n_k} - y_{n_k} \rangle. \tag{45}$$

Thus, taking limit as $k \rightarrow \infty$, we obtain that

$$\langle \bar{u}, x^* - y^* \rangle \leq 0. \tag{46}$$

On the hand, since $x^* \notin \Gamma$, we have from Lemma 2.10 that $x^* \neq y^*$. Hence, we get

$$\langle \bar{u}, x^* - y^* \rangle = \frac{1}{\rho} \langle y^* - (x^* - \rho \bar{u}) + (x^* - y^*), x^* - y^* \rangle > \frac{1}{\rho} \langle x^* - y^*, x^* - y^* \rangle > 0, \tag{47}$$

which is a contradiction to Eq. 46. Therefore, $\liminf_{k \rightarrow \infty} \alpha_{n_k} > 0$.

(b) From (a), we have that $\liminf_{k \rightarrow \infty} \alpha_{n_k} > 0$. Thus, we obtain from Eq. 43 that

$$\begin{aligned} 0 &\leq \limsup_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\|^2 \leq \limsup_{k \rightarrow \infty} (\alpha_{n_k} \|v_{n_k} - y_{n_k}\|^2) \left(\limsup_{k \rightarrow \infty} \frac{1}{\alpha_{n_k}} \right) \\ &= \left(\limsup_{k \rightarrow \infty} \alpha_{n_k} \|v_{n_k} - y_{n_k}\|^2 \right) \left(\frac{1}{\liminf_{k \rightarrow \infty} \alpha_{n_k}} \right) \\ &= 0. \end{aligned}$$

Therefore, we obtain that

$$\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0.$$

□

Now, we are in position to give the main theorem of this section.

Theorem 4.12 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.2. Then, under assumption (8) and Assumption 3.1, we have that $\{x_n\}$ converges to an element of Γ .*

Proof By Lemma 4.3, $\{x_n\}$ is bounded. Thus, there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $\{x_{n_k}\}$ converges to some point x^* . Also, we have that

$$\|v_{n_k} - x_{n_k}\| = \theta_{n_k} \|x_{n_k} - x_{n_k-1}\| \rightarrow 0, \text{ as } k \rightarrow \infty. \tag{48}$$

We now claim that $x^* \in \Gamma$.

Suppose on the contrary that $x^* \notin \Gamma$. Then, it follows from Lemma 4.11 (b) and Eq. 48 that

$$\lim_{k \rightarrow \infty} y_{n_k} = \lim_{k \rightarrow \infty} P_C(v_{n_k} - \rho_{n_k} u_{n_k}) = \lim_{k \rightarrow \infty} x_{n_k} = x^*. \tag{49}$$

Now, without loss of generality, we may assume that $\rho_{n_k} \rightarrow \rho^*$ and $u_{n_k} \rightarrow u^*$. Since F is continuous, it is outer-semicontinuous. Thus, we obtain that $u^* \in F(x^*)$. Therefore, we obtain from Eq. 49 that

$$P_C(x^* - \rho^* u^*) = x^*.$$

It then follows from Lemma 2.10 that $x^* \in \Gamma$, which is a contraction. Hence, our claim holds.

We now show that $\{x_n\}$ converges to x^* .

Replacing z by x^* in Lemma 4.3, we obtain that $\lim_{n \rightarrow \infty} \|x_n - x^*\|^2$ exists. Since x^* is an accumulation point of $\{x_n\}$, we obtain that $\{x_n\}$ converges to x^* . □

Remark 4.13 In view of Corollaries 4.5-4.10, we can obtain various corollaries of Theorem 4.12. Furthermore, in the case that $\theta_n = 0$ for all $n \geq 1$, assumptions (8) and (14) are automatically satisfied. Moreover, we have in this case that $t_n = 1$ for all $n \geq 1$. Hence, we can employ **Procedure A** (see page 1) to obtain similar result as in He et al. (2019, Theorem 3.1).

Algorithm 4.1.

- Step 0:** Let $x_1 \in \mathbb{R}^N$ be given arbitrary and fix $\gamma, \sigma \in (0, 1), 0 < \rho_0 \leq \rho_1 < \infty$. Set $C_1 = \mathbb{R}^N, \bar{x}_1 = x_1$ and $n = 1$.
- Step 1.** Apply **Procedure A** to obtain $x_n = R(\bar{x}_n)$.
- Step 2.** Choose $u_n \in F(x_n)$ and $\rho_n \in [\rho_0, \rho_1]$. Then, compute $y_n = P_C(x_n - \rho_n u_n)$. If $x_n = y_n$: STOP. Otherwise, go to **Step 2**.

Step 3. Compute

$$z_n = \alpha_n y_n + (1 - \alpha_n)x_n$$

and choose the largest $\alpha \in \{\gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists a point $w_n \in F(z_n)$ satisfying

$$\langle w_n, x_n - y_n \rangle \geq \sigma \langle u_n, x_n - y_n \rangle. \tag{50}$$

Step 4. Set $C_n = \{y \in \mathbb{R}^N : \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \cap_{i=1}^n C_i$. Then, compute

$$\bar{x}_{n+1} = P_{C_n^*}(x_n).$$

If $\bar{x}_{n+1} = x_n$, then stop. Otherwise, let $n = n + 1$ and return to Step 1.

Corollary 4.14 (see for example, He et al. (2019, Theorem 3.1)) *Let $\{x_n\}$ be a sequence generated by Algorithm 4.1 such that the following assumptions hold:*

- (a) *The set C is described as in **procedure A** (see page 1).*
- (b) *$F : C \rightrightarrows \mathbb{R}^N$ is locally bounded and continuous.*
- (c) *Γ is nonempty and satisfies condition (7).*

Then, we have that $\{x_n\}$ converges to an element of Γ .

Proof It follows carefully from Lemma 2.13 and Theorem 4.12. □

Remark 4.15 Under the settings of Remark 4.13, we can obtain in general, similar result as in He et al. (2019, Theorem 3.2) without **Procedure A**.

Algorithm 4.2.

Step 0: Let $x_1 \in C$ be given arbitrary and fix $\gamma, \sigma \in (0, 1), 0 < \rho_0 \leq \rho_1 < \infty$. Set $C_1 = \mathbb{R}^N$ and $n = 1$.

Step 1. Choose $u_n \in F(x_n)$ and $\rho_n \in [\rho_0, \rho_1]$. Then, compute $y_n = P_C(x_n - \rho_n u_n)$. If $x_n = y_n$: STOP. Otherwise, go to **Step 2**.

Step 2. Compute

$$z_n = \alpha_n y_n + (1 - \alpha_n)x_n$$

and choose the largest $\alpha \in \{\gamma, \gamma^2, \gamma^3, \dots\}$ such that there exists a point $w_n \in F(z_n)$ satisfying

$$\langle w_n, x_n - y_n \rangle \geq \sigma \langle u_n, x_n - y_n \rangle. \tag{51}$$

Step 3. Set $C_n = \{y \in \mathbb{R}^N : \langle w_n, y - z_n \rangle \leq 0\}$ for $n \geq 2$ and $C_n^* = \cap_{i=1}^n C_i$. Then, compute

$$x_{n+1} = P_C \cap C_n^*(x_n).$$

If $x_{n+1} = x_n$, then stop. Otherwise, let $n = n + 1$ and return to Step 1.

Corollary 4.16 (see, for example, He et al. (2019, Theorem 3.2)) *Let $\{x_n\}$ be a sequence generated by Algorithm 4.2 such that the following assumptions hold:*

- (a) *The feasible set C is a nonempty closed and convex subset of \mathbb{R}^N .*
- (b) *$F : C \rightrightarrows \mathbb{R}^N$ is locally bounded and continuous.*
- (c) *Γ is nonempty and satisfies condition (7).*

Then, we have that $\{x_n\}$ converges to an element of Γ .

Proof It follows directly from Corollary 4.14. □

5 Convergence Analysis for Algorithm 3.3

Remark 5.1 Notice that Step 2 (the linesearch procedure) of Algorithm 3.2 is not utilized in the proof of Lemma 4.1-Lemma 4.3. Thus, Lemma 4.1-Lemma 4.3 hold automatically if $\{x_n\}$ is generated by Algorithm 3.3. Therefore, we only need to prove the version of Lemma 4.11 and Theorem 4.12 corresponding to Algorithm 3.3 in this section.

Lemma 5.2 *Let the sequence $\{x_n\}$ be generated by Algorithm 3.3 such that assumption (8) and Assumption 3.1 are satisfied. Then, we have*

- (a) $\lim_{n \rightarrow \infty} \alpha_n \|y_n - v_n\|^2 = 0$.
- (b) *If there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $\{x_{n_k}\}$ converges to x^* , then $\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0$.*

Proof (a) From Eq. 4, Step 2 and the fact that $x_{n+1} \in C_n^*$, we obtain that

$$\begin{aligned}
 \alpha_n \|v_n - y_n\|^2 &\leq \frac{2\alpha_n}{\sigma} \langle w_n, v_n - y_n \rangle \\
 &\leq \frac{2}{\sigma} \langle w_n, v_n - z_n \rangle \\
 &\leq \frac{2}{\sigma} (\langle w_n, v_n - x_{n+1} \rangle + \langle w_n, x_{n+1} - z_n \rangle) \\
 &\leq \frac{2}{\sigma} \|w_n\| \|v_n - x_{n+1}\|.
 \end{aligned}
 \tag{52}$$

Since $\{z_n\}$ is bounded and F is locally bounded, we obtain from Proposition 2.3 that $\{w_n\}$ is also bounded. Thus, we obtain from Eq. 52 and Lemma 4.3 that

$$\lim_{n \rightarrow \infty} \alpha_n \|y_n - v_n\|^2 = 0.
 \tag{53}$$

- (b) Since $\{\alpha_n\} \subset (0, 1)$ is bounded, we have that $\liminf_{n \rightarrow \infty} \alpha_n \geq 0$.

We now consider two possible cases:

Case 1: Suppose that $\liminf_{n \rightarrow \infty} \alpha_n = 0$. Then, we can choose a subsequence of $\{\alpha_n\}$ denoted by $\{\alpha_{n_k}\}$ such that $\lim_{k \rightarrow \infty} \alpha_{n_k} = 0$ and

$$\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = t \geq 0. \tag{54}$$

Now, define $\bar{\alpha}_{n_k} := \frac{\alpha_{n_k}}{\gamma}$. Then, $\bar{z}_{n_k} := \bar{\alpha}_{n_k} y_{n_k} + (1 - \bar{\alpha}_{n_k}) v_{n_k}$. Since $\alpha_{n_k} \rightarrow 0$ as $k \rightarrow \infty$, we obtain that $\bar{\alpha}_{n_k} \rightarrow 0$ as $k \rightarrow \infty$. Hence,

$$\lim_{k \rightarrow \infty} \|\bar{z}_{n_k} - v_{n_k}\| = 0. \tag{55}$$

Now, from the definition of \bar{z}_{n_k} and Step 2, we obtain that

$$\langle \bar{w}_{n_k}, v_{n_k} - y_{n_k} \rangle < \frac{\sigma}{2} \|v_{n_k} - y_{n_k}\|^2,$$

which implies that

$$2\langle \bar{w}_{n_k} - u_{n_k}, v_{n_k} - y_{n_k} \rangle + 2\langle u_{n_k}, v_{n_k} - y_{n_k} \rangle < \sigma \|v_{n_k} - y_{n_k}\|^2. \tag{56}$$

Now, set $s_{n_k} := v_{n_k} - \rho_{n_k} u_{n_k}$. Then, Eq. 56 becomes

$$2\langle \bar{w}_{n_k} - u_{n_k}, v_{n_k} - y_{n_k} \rangle + \frac{2}{\rho_{n_k}} \langle v_{n_k} - s_{n_k}, v_{n_k} - y_{n_k} \rangle < \sigma \|v_{n_k} - y_{n_k}\|^2,$$

which implies that

$$2\langle \bar{w}_{n_k} - u_{n_k}, v_{n_k} - y_{n_k} \rangle + \frac{1}{\rho_{n_k}} \left(\|v_{n_k} - y_{n_k}\|^2 + \|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \right) < \sigma \|v_{n_k} - y_{n_k}\|^2.$$

That is,

$$\frac{1}{\rho_{n_k}} \left(\|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \right) < \left(\sigma - \frac{1}{\rho_{n_k}} \right) \|v_{n_k} - y_{n_k}\|^2 - 2\langle \bar{w}_{n_k} - u_{n_k}, v_{n_k} - y_{n_k} \rangle. \tag{57}$$

Now, by Lemma 4.2, we obtain that $\lim_{k \rightarrow \infty} \|x_{n_k} - v_{n_k}\| = 0$. Thus, since $x_{n_k} \rightarrow x^*$, we have that $v_{n_k} \rightarrow x^*$. Using Assumption 3.1 (b), the boundedness of $\{v_{n_k}\}$ and Proposition 2.3, we obtain that $\{u_{n_k}\}$ is also bounded. Thus, we can choose a subsequence of $\{u_{n_k}\}$ still denoted by $\{u_{n_k}\}$ such that $u_{n_k} \rightarrow \bar{u}$. Since F is continuous, it is outer-semicontinuous. Hence, $\bar{u} \in F(x^*)$. We also assume without loss of generality that $\rho_{n_k} \rightarrow \rho \in [\rho_0, \rho_1] \subset [\rho_0, \frac{1}{\sigma})$. Again, from Eq. 55, we obtain that $\bar{z}_{n_k} \rightarrow x^*$. Since F is inner-semicontinuous and $\bar{u} \in F(x^*)$, we can choose a subsequence $w_{n_k} \in F(\bar{z}_{n_k})$ such that $\bar{w}_{n_k} \rightarrow \bar{u}$.

Also, since $\{v_{n_k}\}$, $\{u_{n_k}\}$, $\{y_{n_k}\}$ and $\{\bar{w}_{n_k}\}$ are bounded, we can choose a subsequence $\{k_j\}$ of $\{k\}$ such that

$$\begin{aligned} & \frac{1}{\rho} \left[\limsup_{k \rightarrow \infty} \left(\|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \right) \right] \\ & \leq \limsup_{k \rightarrow \infty} \left[\left(\sigma - \frac{1}{\rho_{n_k}} \right) \|v_{n_k} - y_{n_k}\|^2 - 2 \langle \bar{w}_{n_k} - u_{n_k}, v_{n_k} - y_{n_k} \rangle \right] \\ & = \lim_{j \rightarrow \infty} \left[\left(\sigma - \frac{1}{\rho_{n_{k_j}}} \right) \|v_{n_{k_j}} - y_{n_{k_j}}\|^2 - 2 \langle \bar{w}_{n_{k_j}} - u_{n_{k_j}}, v_{n_{k_j}} - y_{n_{k_j}} \rangle \right]. \end{aligned}$$

Thus, we obtain from Eq. 54 that

$$\limsup_{k \rightarrow \infty} \left(\|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \right) \leq \rho \left(\sigma - \frac{1}{\rho} \right) t. \tag{58}$$

At this point, we claim that $t = 0$. Otherwise, Eq. 58 will become

$$\limsup_{k \rightarrow \infty} \left(\|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \right) \leq \rho \left(\sigma - \frac{1}{\rho} \right) t < 0.$$

But for $\varepsilon = \frac{-\rho \left(\sigma - \frac{1}{\rho} \right)}{2} t > 0$, there exists $N \in \mathbb{N}$ such that

$$\|s_{n_k} - v_{n_k}\|^2 - \|s_{n_k} - y_{n_k}\|^2 \leq \rho \left(\sigma - \frac{1}{\rho} \right) + \varepsilon = \frac{\rho \left(\sigma - \frac{1}{\rho} \right)}{2} < 0 \quad \forall k \in \mathbb{N}, \quad k \geq N.$$

Thus, we obtain that

$$\|v_{n_k} - s_{n_k}\| < \|y_{n_k} - s_{n_k}\| \quad \forall k \in \mathbb{N},$$

which is a contradiction to the definition of $y_{n_k} = P_C(v_{n_k} - \rho_{n_k} u_{n_k})$. Therefore, $t = 0$. Hence, Eq. 54 becomes

$$\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0.$$

Case 2: Suppose that $\liminf_{n \rightarrow \infty} \alpha_n > 0$. Then, we obtain from Eq. 53 that

$$\begin{aligned} 0 & \leq \limsup_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\|^2 \leq \limsup_{k \rightarrow \infty} (\alpha_{n_k} \|v_{n_k} - y_{n_k}\|^2) \left(\limsup_{k \rightarrow \infty} \frac{1}{\alpha_{n_k}} \right) \\ & = \left(\limsup_{k \rightarrow \infty} \alpha_{n_k} \|v_{n_k} - y_{n_k}\|^2 \right) \left(\frac{1}{\liminf_{k \rightarrow \infty} \alpha_{n_k}} \right) \\ & = 0. \end{aligned}$$

Therefore, we obtain that

$$\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0.$$

□

Theorem 5.3 *Let $\{x_n\}$ be a sequence generated by Algorithm 3.3. Then, under assumption (8) and Assumption 3.1, we have that $\{x_n\}$ converges to an element of Γ .*

Proof By Lemma 4.3, $\{x_n\}$ is bounded. Thus, there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $\{x_{n_k}\}$ converges to some point x^* . Thus, we obtain from Lemma 5.2 (b) that

$$\lim_{k \rightarrow \infty} \|v_{n_k} - y_{n_k}\| = 0. \tag{59}$$

Also, from Lemma 4.2, we obtain that

$$\lim_{k \rightarrow \infty} \|v_{n_k} - x_{n_k}\| = 0. \tag{60}$$

Hence, from Eqs. 59 and 60, we obtain

$$\lim_{k \rightarrow \infty} y_{n_k} = \lim_{k \rightarrow \infty} P_C(v_{n_k} - \rho_{n_k} u_{n_k}) = \lim_{k \rightarrow \infty} x_{n_k} = x^*. \tag{61}$$

Now, without loss of generality, we may assume that $\rho_{n_k} \rightarrow \rho^*$ and $u_{n_k} \rightarrow u^*$. Since F is continuous, it is outer-semicontinuous. Thus, we obtain that $u^* \in F(x^*)$. Therefore, we obtain from Eq. 61 that

$$P_C(x^* - \rho^* u^*) = x^*.$$

It then follows from Lemma 2.10 that $x^* \in \Gamma$.

We now show that $\{x_n\}$ converges to x^* .

Replacing z by x^* in Lemma 4.3, we obtain that $\lim_{n \rightarrow \infty} \|x_n - x^*\|^2$ exists. Since x^* is an accumulation point of $\{x_n\}$, we obtain that $\{x_n\}$ converges to x^* . □

Remark 5.4 Following Remark 4.13, we can also obtain various corollaries of Theorem 5.3.

6 Numerical Experiments

In this section, we discuss the numerical behavior of Algorithm 3.2 and Algorithm 3.3 using test examples taken from the literature. We only compare our methods with Algorithms 1.3 and 1.4 of He et al. (2019) since in He et al. (2019, Section 4), we have that the methods in He et al. (2019) are more efficient than most relevant methods in the literature.

The codes are implemented in Matlab 2016 (b). We perform all computations on a personal computer with an Intel(R) Core(TM) i5-2600 CPU at 2.30GHz and 8.00 Gb-RAM.

We consider the same set of examples considered in He et al. (2019, Section 4). We randomly choose $x_0, x_1 \in \mathbb{R}^N$ and the inertial factor θ_n satisfying assumptions (8) and (14).

Example 6.1 Consider the following convex non-smooth optimization problem (see also Dong et al. 2017; He et al. 2019)

$$\min_{x \in C} \varphi(x),$$

where $\varphi(x) = -x_1 + 20 \max\{x_1^2 + x_2^2 - 1, 0\}$ and $C = \{x \in \mathbb{R}_+^2 : x_1 + x_2 \leq 1\}$. This problem is equivalent to the MVIP (1) with $F(x) = \partial\varphi(x)$, where $\partial\varphi(x)$ is the subdifferential of φ at x :

$$\partial\varphi(x) = \begin{cases} (-1 + 40x_1, 40x_2), & \text{if } \|x\| > 1; \\ (-1, 0), & \text{if } \|x\| < 1; \\ \{(-1 + 40tx_1, 40tx_2) | t \in [0, 1]\}, & \text{if } \|x\| = 1. \end{cases}$$

Then, we see that $x^* = (1, 0)$ is the unique solution of the problem, and the multivalued mapping $F = \partial\varphi$ satisfies the assumptions of Assumption 3.1 (b).

For the parameters, we choose $\rho_n \in (0, 2)$, $\sigma = 0.8$, and $\gamma = 0.7$. Furthermore, we take $\|x_n - x^*\| \leq \epsilon$ as the termination criterion. We stress that these choices are the same as the ones considered by He et al. (2019) for their numerical experiments.

For $\epsilon = 10^{-7}$, we obtain the numerical results listed in Table 1 and Fig. 1, which show that our methods perform better than Algorithm 1.3 and Algorithm 1.4 of He et al. (2019).

For $\epsilon = 10^{-10}$, it was observed in He et al. (2019, Section 4) that Algorithm 1.3 of He et al. (2019) does not work well because of the presence of **Procedure A** in the iterative steps. Therefore, in this setting, we shall compare our methods with only Algorithm 1.4 of He et al. (2019). For this, we obtain the numerical results reported in Table 2 and Fig. 2, which show that our methods still perform better than Algorithm 1.4 of He et al. (2019).

We consider the following cases for the numerical experiments of Example 6.1.

- Case 1:** $x_1 = (0.5, -0.25)^T, x_0 = (0.5, -0.25)^T$ and $\theta_n = \frac{2n-1}{8n}$.
- Case 2:** $x_1 = (0.7, 0.25)^T, x_0 = (0.5, 0.25)^T$ and $\theta_n = \frac{2n-1}{8n}$.
- Case 3:** $x_1 = (-1.5, 1)^T, x_0 = (1, -0.2)^T$ and $\theta_n = \frac{n-1}{n+4}$.
- Case 4:** $x_1 = (-0.5, 1.5)^T, x_0 = (-0.5, 1)^T$ and $\theta_n = \frac{n-1}{n+4}$.

Table 1 Numerical results for Example 6.1 with $\epsilon = 10^{-7}$

		Alg. 3.2	Alg. 3.3	Alg. 1.3	Alg. 1.4
Case 1	CPU time (sec) No. of Iteration	0.0720 5	0.0150 3	0.1310 10	0.2550 17
Case 2	CPU time (sec) No. of Iteration	0.0724 5	0.0151 3	0.1330 10	0.2550 17
Case 3	CPU time (sec) No. of Iteration	0.0720 5	0.0350 4	0.1320 10	0.2550 18
Case 4	CPU time (sec) No. of Iteration	0.0480 4	0.0430 4	0.1040 8	0.2610 18




Fig. 1 $\|x_n - x^*\|$ vs Iteration numbers (n) for Example 6.1 with $\epsilon = 10^{-7}$: Top Left: **Case 1**; Top Right: **Case 2**; Bottom Left: **Case 3**; Bottom Right: **Case 4**

Table 2 Numerical results for Example 6.1 with $\epsilon = 10^{-10}$

		Alg. 3.2	Alg. 3.3	Alg. 1.4
Case 1	CPU time (sec) No. of Iteration	0.1420 10	0.0160 4	0.7910 34
Case 2	CPU time (sec) No. of Iteration	0.1390 10	0.0110 4	0.7840 34
Case 3	CPU time (sec) No. of Iteration	0.1010 8	0.0370 5	0.7810 34
Case 4	CPU time (sec) No. of Iteration	0.1020 8	0.0370 5	0.7920 35




Fig. 2 $\|x_n - x^*\|$ vs Iteration numbers (n) for Example 6.1 with $\epsilon = 10^{-10}$: Top Left: **Case 1**; Top Right: **Case 2**; Bottom Left: **Case 3**; Bottom Right: **Case 4**

Example 6.2 We next consider the following optimization problem which was also considered in He et al. (2019) and Ye and He (2015).

$$\min_{x \in C} \varphi(x),$$

where $C = \left\{ x \in \mathbb{R}^5 : x_i \geq 0, i = 1, 2, \dots, 5, \sum_{i=1}^5 x_i = a, a > 0 \right\}$ and $\varphi(x) = \frac{0.5(Hx, x) + (q, x) + 1}{\sum_{i=1}^5 x_i}$.

Furthermore, H denotes a positive diagonal matrix with the same element h taken from the interval $(0.1, 2)$ and $q = (-1, -1, -1, -1, -1)$. Clearly, this problem

Table 3 Numerical results for Example 6.2 with $\epsilon = 10^{-4}$

		Alg. 3.2	Alg. 3.3	Alg. 1.4
Case 1	CPU time (sec) No. of Iteration	0.5210 14	0.0610 6	1.1210 42
Case 2	CPU time (sec) No. of Iteration	0.3100 10	0.0830 4	0.9220 33
Case 3	CPU time (sec) No. of Iteration	0.2200 9	0.0800 4	0.9210 32
Case 4	CPU time (sec) No. of Iteration	0.2920 9	0.0870 4	0.9290 32




Fig. 3 $\|x_n - x^*\|$ vs Iteration numbers (n) for Example 6.2 with $\epsilon = 10^{-4}$: Top Left: **Case 1**; Top Right: **Case 2**; Bottom Left: **Case 3**; Bottom Right: **Case 4**

is equivalent to MVIP (1) with solution set $\Gamma = \{\frac{1}{5}(a, \dots, a)\}$, where $F(x) = (\varphi_1(x), \dots, \varphi_5(x))$ and

$$\varphi_i(x) = \frac{\partial \varphi(x)}{\partial x_i} = \frac{hx_i \sum_{i=1}^5 x_i - 0.5h \sum_{i=1}^5 x_i^2 - 1}{\left(\sum_{i=1}^5 x_i\right)^2}.$$

For $\epsilon = 10^{-4}$, $\sigma = 0.3$ and for some randomly chosen values of a , we compare our methods with Algorithm 1.4 of He et al. (2019). We obtain the numerical results displayed in Table 3 and Fig. 3, which show that our methods perform better than Algorithm 1.4 of He et al. (2019).

We consider the following cases for the numerical experiments of Example 6.2.

- Case 1:** $x_1 = (1, 0.5, 1, 1.5, 1)^T$, $x_0 = (1, 0.5, 1, 1.5, 1)^T$, $a = 5$ and $\theta_n = \frac{2n-1}{8n}$.
- Case 2:** $x_1 = (3, 2, 2, 1, 2)^T$, $x_0 = (4.3, 2.5, 2.2, 0.3, 0.7)^T$, $a = 10$ and $\theta_n = \frac{2n-1}{8n}$.
- Case 3:** $x_1 = (0.1, 0.9, 2, 0.5, 1.5)^T$, $x_0 = (0.3, 0.5, 1.2, 2.5, 0.5)^T$, $a = 5$ and $\theta_n = \frac{n-1}{n+4}$.
- Case 4:** $x_1 = (2.1, 2.9, 2, 1.5, 1.5)^T$, $x_0 = (1.3, 1.5, 2.2, 3.5, 1.5)^T$, $a = 10$ and $\theta_n = \frac{n-1}{n+4}$.

7 Conclusion

We propose two new inertial extrapolation projection-type methods for solving MVIPs when the multivalued mapping F is only required to be locally bounded without any monotonicity assumption. The first method uses a linesearch as in He et al. (2019, Algorithms 1 and 2) while the second method uses a different linesearch procedure with the aim of minimizing the number of evaluation of the multivalued mapping F in each search. Furthermore, our inertial techniques for establishing the convergence of these methods are quite different from the commonly used ones in most papers (see for example Chalamjiak et al. 2018; Chuang 2017; Ochs et al. 2015; Lorenz and Pock 2015; Polyak 1964; Shehu and Chalamjiak 2019; Lorenz and Pock 2015; Mainge 2008; Moudafi and Oliny 2003; Shehu et al. 2019; Shehu et al. 2019; Thong and Hieu 2018; Thong and Hieu 2017 and the references therein). Moreover, based on the weaker assumptions on the inertial factor in our methods, we derive several special cases of our methods. Finally, we consider some numerical implementations of our methods and compare them with the methods in He et al. (2019, Algorithms 1 and 2), in order to show the profits that we gain by introducing the new inertial extrapolation steps. In fact, in all our comparisons, the numerical results demonstrate that our methods perform better than the methods in He et al. (2019, Algorithms 1 and 2). Thus, our results improve and generalize many recent important results in this direction.

Acknowledgements The authors sincerely thank the Editor-in-Chief and anonymous referees for their careful reading, constructive comments and fruitful suggestions that help improve the manuscript. The research of the first author is supported by the National Research Foundation (NRF) South Africa (S& F-DSI/NRF Free Standing Postdoctoral Fellowship; Grant Number: 120784). The first author also acknowledges the financial support from DSI/NRF, South Africa Center of Excellence in Mathematical and Statistical Sciences (CoE-MaSS) Postdoctoral Fellowship. The second author has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Program (FP7 - 2007-2013) (Grant agreement No. 616160).

Funding Open Access funding provided by Institute of Science and Technology (IST Austria).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alvarez F, Attouch H (2001) An inertial proximal method for maximal monotone operators via discretization of a nonlinear oscillator with damping. *Set-Valued Anal* 9:3–11
- Attouch H, Cabot A (2019) Convergence of a relaxed inertial proximal algorithm for maximally monotone operators. *Math Program*. <https://doi.org/10.1007/s10107-019-01412-0>
- Attouch H, Cabot A (2019) Convergence of a relaxed inertial forward–backward algorithm for structured monotone inclusions. *Appl Math Optim* 80:547–598
- Attouch H, Goudon X, Redont P (2000) The heavy ball with friction. I. The continuous dynamical system. *Commun Contemp Math* 2(1):1–34
- Beck A, Teboulle M (2009) A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J Imaging Sci* 2(1):183–202
- Bot RI, Csetnek ER, Hendrich C (2015) Inertial Douglas-Rachford splitting for monotone inclusion. *Appl Math Comput* 256:472–487
- Bot RI, Csetnek ER (2016) An inertial alternating direction method of multipliers. *Minimax Theory Appl* 1:29–49
- Brouwer LEJ (1912) Über Abbildung von Mannigfaltigkeiten. *Math Ann* 71(4):97–115
- Browder FE (1965) Multi-valued monotone nonlinear mappings and duality mappings in Banach spaces. *Trans Am Math Soc* 18:338–351
- Carey M, Ge YE (2012) Comparison of methods for path flow reassignment for dynamic user equilibrium. *Netw Spat Econ* 12:337–376
- Censor Y, Gibali A, Reich S (2011) The subgradient extragradient method for solving variational inequalities in Hilbert space. *J Optim Theory Appl* 148:318–335
- Chambolle A, Dossal C. h. (2015) On the convergence of the iterates of the “fast iterative shrinkage/thresholding algorithm”. *J Optim Theory Appl* 166:968–982
- Cholamjiak W, Cholamjiak P, Suantai S (2018) An inertial forward-backward splitting method for solving inclusion problems in Hilbert spaces. *J Fixed Point Theory Appl*, 20. <https://doi.org/10.1007/s11784-018-0526-5>
- Chuang CS (2017) Hybrid inertial proximal algorithm for the split variational inclusion problem in Hilbert spaces with applications. *Optimization* 66(5):777–792
- Dong QL, Lu YY, Yang J, He S (2017) Approximately solving multi-valued variational inequalities by using a projection and contraction algorithm. *Numer Algor* 76:799–812

- Facchinei F, Pang JS (2003) Finite-dimensional variational inequalities and complementarity problems vol 1 & 2. Springer, Berlin
- Fang CJ, Chen SL (2014) A subgradient extragradient algorithm for solving multi-valued variational inequality. *Appl Math Comput* 229:123–130
- Goebel K, Reich S (1984) Uniform convexity, hyperbolic geometry, and nonexpansive mappings. Marcel Dekker, New York
- He YR (2006) A new double projection algorithm for variational inequalities. *J Comput Appl Math* 185:66–173
- He X, Huang N, Li X (2019) Modified projection methods for solving multi-valued variational inequality without monotonicity. *Netw Spat Econ*. <https://doi.org/10.1007/s11067-019-09485-2>
- Konnov IV (1998) A combined relaxation method for variational inequalities with nonlinear constraints. *Math Program* 80:239–252
- Korpelevich GM (1976) The extragradient method for finding saddle points and other problems. *Matecon* 12:747–756
- Lorenz DA, Pock T (2015) An inertial forward–backward algorithm for monotone inclusions. *J Math Imaging Vis* 51:311–325
- Mainge PE (2008) Convergence theorems for inertial KM-type algorithms. *J Comput Appl Math* 219(1):223–236
- Moudafi A, Oliny M (2003) Convergence of a splitting inertial proximal method for monotone operators. *J Comput Appl Math* 155:447–454
- Nadler SB (1969) Multi-valued contraction mappings. *Pac J Math (English Series)* 30:475–488
- Nesterov Y (1983) A method of solving a convex programming problem with convergence rate $O(1/k^2)$. *Soviet Math Doklady* 27:372–376
- Ochs P, Brox T, Pock T (2015) iPiasco: inertial proximal algorithm for strongly convex optimization. *J Math Imaging Vis* 53:171–181
- Oggioni G, Smeers Y, Allevi E, Schaible S (2012) A generalized Nash equilibrium model of market coupling in the European power system. *Netw Spat Econ* 12:503–560
- Polyak BT (1964) Some methods of speeding up the convergence of iterates methods. *USSR Comput Math Phys* 4(5):1–17
- Raciti F, Falsaperla P (2007) Improved noniterative algorithm for the calculation of the equilibrium in the traffic network problem. *J Optim Theory Appl* 133:401–411
- Rockafellar RT, Wets JB (2004) Variational analysis. Springer, New York
- Rockafellar RT (1970) Convex analysis. Princeton University Press, Princeton
- Shehu Y, Cholamjiak P (2019) Iterative method with inertial for variational inequalities in Hilbert spaces. *Calcolo*, 56(1)
- Shehu Y, Li XH, Dong QL (2019) An efficient projection-type method for monotone variational inequalities in Hilbert spaces. *Numer Algorithms*. <https://doi.org/10.1007/s11075-019-00758-y>
- Shehu Y, Vuong PT, Zemkoho A (2019) An inertial extrapolation method for convex simple bilevel optimization. *Optim Methods Softw*. <https://doi.org/10.1080/10556788.2019.1619729>
- Thong DV, Hieu DV (2018) Inertial subgradient extragradient algorithms with line-search process for solving variational inequality problems and fixed point problems. *Numer Algorithms*. <https://doi.org/10.1007/s11075-018-0527-x>
- Thong DV, Hieu DV (2017) An inertial method for solving split common fixed point problems. *J Fixed Point Theory Appl* 19(4):3029–3051
- Tseng P (2000) A modified forward-backward splitting method for maximal monotone mappings. *SIAM J Control Optim* 38:431–446
- Vuong PT (2019) The global exponential stability of a dynamical system for solving variational inequalities. *Netw Spat Econ*. <https://doi.org/10.1007/s11067-019-09457-6>
- Ye ML, He YR (2015) A double projection method for solving variational inequalities without monotonicity. *Comput Optim Appl* 60:141–150